arXiv:math/0601208v2 [math.DG] 1 Aug 2006

EXISTENCE AND UNIQUENESS FOR P-AREA MINIMIZERS IN THE HEISENBERG GROUP

JIH-HSIN CHENG, JENN-FANG HWANG, AND PAUL YANG

Abstract. In [3], we studied p-mean curvature and the associated p-minimal surfaces in the Heisenberg group from the viewpoint of PDE and differential geometry. In this paper, we look into the problem through the variational formulation. We study a generalized p-area and associated (p-) minimizers in general dimensions.

We prove the existence and investigate the uniqueness of minimizers. Since this is reduced to solving a degenerate elliptic equation, we need to consider the effect of the singular set and this requires a careful study. We define the notion of weak solution and prove that in a certain Sobolev space, a weak solution is a minimizer and vice versa. We also give many interesting examples in dimension 2. An intriguing point is that, in dimension 2, a C^2-smooth solution from the PDE viewpoint may not be a minimizer. However, this statement is true for higher dimensions due to the relative smallness of the size of the singular set.

1. Introduction and statement of the results

The p-minimal (or X-minimal, H-minimal in the terminology of some authors, e.g., [6], [7], [16]) surfaces have been studied extensively in the framework of geometric measure theory. Starting from the work [3], we studied the subject from the viewpoint of partial differential equations and that of differential geometry (we use the term p-minimal since this is the notion of minimal surfaces in pseudohermitian geometry; "p" stands for "pseudohermitian").

Let Ω be a bounded domain in R^{2n}. Let $\vec{X} = (x_1, x_{1'}, x_2, x_{2'}, .., x_n, x_{n'}) \in \Omega$. For a graph $(\vec{X}, u(\vec{X}))$ in the Heisenberg group of dimension $2n+1$ with prescribed p-mean curvature $H = H(\vec{X})$, the equation for $u : \Omega \subset R^{2n} \to R$ reads

$$div \frac{\nabla u - \vec{X}^*}{|\nabla u - \vec{X}^*|} = H \tag{1.1}$$

where $\vec{X}^* = (x_{1'}, -x_1, x_{2'}, -x_2, ..., x_{n'}, -x_n)$ (see (2.10) in Section 2 for a geometric interpretation). In general, for a vector field $\vec{G} = (g_1, g_2, ..., g_{2n})$ on $\Omega \subset R^{2n}$, we define $\vec{G}^* \equiv (g_2, -g_1, g_4, -g_3, ..., g_{2n}, -g_{2n-1})$. The equation (1.1) is the Euler-Lagrange equation (away from the singular set $\nabla u - \vec{X}^* = 0$) of the following energy functional (called the p-area of the graph defined by u if $H = 0$, see Section 2):

$$\mathcal{X}(u) = \int_\Omega \{|\nabla u - \vec{X}^*| + Hu\} dx_1 \wedge dx_{1'} \wedge ... \wedge dx_n \wedge dx_{n'}. \tag{1.2}$$

1991 *Mathematics Subject Classification.* Primary: 35L80; Secondary: 35J70, 32V20, 53A10, 49Q10.

Key words and phrases. Minimizer, p-area, Heisenberg group.

Since we consider the variation over the whole domain, the singular set will cause the main difficulty in the study. In order to explain this, we generalize $\mathcal{X}(\cdot)$ by considering an arbitrary vector field $\vec{F} = \vec{F}(\vec{X})$ instead of $-\vec{X}^*$ in the following form:

$$\mathcal{F}_q(u) \equiv \int_\Omega \{|\nabla u + \vec{F}|^q + qHu\} dx_1 \wedge dx_2 \wedge ... \wedge dx_m \tag{1.3}$$

for $1 \leq q < \infty$, where $\Omega \subset R^m$. Let $S(u)$ denote the singular set of u, consisting of the points where $\nabla u + \vec{F} = 0$. Let $u_\varepsilon = u + \varepsilon\varphi$. It is easy to compute (see Section 3 for the case $q = 1$) the first variation of $\mathcal{F}_q$: (omitting the Euclidean volume element)

$$\begin{aligned} &\frac{d\mathcal{F}_q(u_\varepsilon)}{d\varepsilon}|_{\varepsilon=0\pm} \\ = \ &c_q \int_{S(u)} |\nabla\varphi|^q + \int_{\Omega\setminus S(u)} q|\nabla u + \vec{F}|^{q-2}(\nabla u + \vec{F}) \cdot \nabla\varphi + \int_\Omega qH\varphi \end{aligned} \tag{1.4}$$

where $c_q = \pm 1$ for $q = 1$ and $c_q = 0$ for $1 < q < \infty$.

For $q = 1$, can we ignore the term $\pm\int_{S(u)} |\nabla\varphi|$? A recent paper of Balogh answered this question completely. In [1] Balogh studied the size of the singular set $S(u)$ (called the characteristic set in [1]). He showed (Theorem 3.1(2) in [1]) that for $\vec{F} = -\vec{X}^*$ in R^{2n}, $S(u)$ has locally finite n-dimensional Hausdorff measure if $u \in C^2$. We obtained the same result as Lemma 5.4 in [3] by a different argument (we used only elementary linear algebra and the implicit function theorem in the proof; also we were not aware of [1] at the time [3] was written). In this paper, we generalize this result to the situation of general $\vec{F}$ (see Theorem D below and its proof in Section 6). For $u \in C^{1,1}$ and $\vec{F} = -\vec{X}^*$ in R^{2n}, Balogh showed (Theorem 3.1(1) in [1]) that $dim_E S(u) < 2n - \delta$ where dim_E denotes the Hausdorff dimension with respect to the Euclidean metric and δ depends on the Lipschitz constant of ∇u. He also proved the existence of $u \in \cap_{0<\alpha<1} C^{1,\alpha}$ such that $S(u)$ has positive Lebesgue measure for any $\vec{F} \in C^1(\Omega)$ where $\Omega \subset R^m$ is a given bounded domain (Theorem 4.1(2) in [1]). In this paper, we consider functions u of class $W^{1,1}$ so that the size of $S(u)$ may be large according to Balogh. Therefore for the case of $q = 1$ in (1.4), we can not neglect the contribution of the singular set to define the weak solutions (see Definition 3.2) to the Euler-Lagrange equation of $\mathcal{F}_q$:

$$div \frac{\nabla u + \vec{F}}{|\nabla u + \vec{F}|^{2-q}} = H. \tag{1.5}$$

Equation (1.5) has been studied in various situations. For $\vec{F} = 0$, $H = 0$, (1.5) is known to be the q-harmonic equation for $1 < q < \infty$ while it is the equation associated to the least gradient problem for $q = 1$ (see, for instance, [19], [12], [11], etc.). Geometrically there is a dichotomy for the 1-form $\Theta \equiv dz + F_I dx_I$ associated to the vector field $\nabla u + \vec{F}$, where $\vec{F} = (F_I)$. Namely, the hyperplane distribution defined by the kernel of Θ might be either integrable or (completely) nonintegrable (Θ is called a contact form in this case). When $\vec{F} = 0$, this is the integrable case. For the nonintegrable case (e.g. $\vec{F} = -\vec{X}^*$), the quantity of the left side in (1.5) with $q = 1$ can be realized as the p-mean curvature of the graph defined by u in pseudohermitian geometry (see (2.12)).

We study equation (1.5) with $q = 1$:

$$div \frac{\nabla u + \vec{F}}{|\nabla u + \vec{F}|} = H \tag{1.6}$$

Definition 1.1. Let Ω be a domain in R^m, $m \geq 1$. We say $u \in C^2(\Omega)$ is a C^2 smooth solution to (1.6) if and only if (1.6) holds in $\Omega \backslash S(u)$.

In [3] and [2], we considered C^2-smooth solutions u to (1.1) (i.e., (1.6) with $\vec{F} = -\vec{X}^*$) with $H = 0$ in dimension 2 and, among other things, we proved a Bernstein-type theorem. Later in [8] the authors obtained a similar Bernstein-type theorem through a different approach. The description of the singular set for a C^2-smooth solution to (1.1) occupies a central position in [3]. As a geometric application, we can show the nonexistence of C^2-smooth, closed surfaces of genus ≥ 2 with bounded p-mean curvature in any pseudohermitian 3-manifold. In [3] we also proved a uniqueness theorem for C^2-smooth solutions for the Dirichlet problem of (1.6) in R^{2n}. Recently Ritoré and Rosales proved a rigidity result for C^2-smooth surfaces of nonzero constant p-mean curvature and an Alexandrov-type theorem in the 3-dimensional Heisenberg group (see Theorem 6.1 and Theorem 6.10 in [18], respectively).

In this paper we consider $W^{1,1}$ minimizers for

$$\mathcal{F}(u) \equiv \int_\Omega \{|\nabla u + \vec{F}| + Hu\} dx_1 \wedge dx_2 \wedge ... \wedge dx_m \tag{1.7}$$

((1.3) with $q = 1$). In Section 3 we define and show that in the space $W^{1,1}$, a minimizer for (1.7) is a weak solution to the equation (1.6) and vice versa (see Theorem 3.3). In order to overcome the trouble caused by singular sets which are not negligible, we introduce the notion of "regular value". Suppose $u \in W^{1,1}$, $\varphi \in W_0^{1,1}$. Define $u_\varepsilon \equiv u + \varepsilon\varphi$ for $\varepsilon \in R$. We prove that there are at most countably many ε's for which

$$\int_{S(u_\varepsilon)} | \nabla \varphi | \neq 0$$

(cf. (1.4) for $q = 1$). We call such an ε singular, otherwise regular. That is, the above integral vanishes for almost all (regular) ε (see Lemma 3.1). So we do not need to worry about the size of the singular set for regular ε's. The idea of considering regular values plays a central role both in the proof of the equivalence between minimizers and weak solutions and in the proof of the uniqueness theorems in Section 5.

In Section 4 we prove the existence of a Lipschitz continuous minimizer for $\mathcal{F}(\cdot)$ with a given boundary value in the case of $H = 0$ under the following condition on $\vec{F}$:

$$\partial_K F_I = \partial_I f_K, \quad I, K = 1, ..., m \tag{1.8}$$

for C^1-smooth functions f_K's (cf. (4.11)). We require Ω to be a p-convex domain (see Definition 4.1).

Theorem A. *Let Ω be a p-convex bounded domain in $R^m, m \geq 2$, with $\partial\Omega \in C^{2,\alpha}$ $(0 < \alpha < 1)$. Let $\varphi \in C^{2,\alpha}(\bar{\Omega})$. Suppose $\vec{F} \in C^{1,\alpha}(\bar{\Omega})$ satisfies the condition (1.8) (or (4.11)) for $C^{1,\alpha}$-smooth and bounded f_K's in Ω. Then there exists a Lipschitz continuous minimizer $u \in C^{0,1}(\bar{\Omega})$ for $\mathcal{F}(\cdot)$ with $H = 0$ such that $u = \varphi$ on $\partial\Omega$.*

We note that a C^2-smooth bounded domain with positively curved (positive principal curvatures) boundary is p-convex. Also condition (1.8) includes the case $\vec{F} = -\vec{X}^*$. We can actually find out all the solutions to (1.8) (see (4.13)). We notice that, for $n = 1$, Pauls ([16]) proved the existence of a continuous $W^{1,p}$ minimizer for $\mathcal{X}(\cdot)$ under the assumption that the graph of the prescribed boundary function φ satisfies the bounded slope condition (see [9]).

The idea of the proof of Theorem A is to invoke Theorem 11.8 in [9] for a family of elliptic approximating equations (see also [16]). Namely we first solve the Dirichlet problem for the following equations:

$$\begin{aligned} Q_\varepsilon u &\equiv div(\frac{\nabla u + \vec{F}}{\sqrt{\varepsilon^2 + |\nabla u + \vec{F}|^2}}) = 0 \quad \text{in } \Omega, \\ u &= \varphi \quad \text{on } \partial\Omega \end{aligned}$$

(see (4.1)). We end up obtaining a uniform C^1 bound for solutions to the above equations, and a subsequence of solutions converges to a Lipschitz continuous minimizer as $\varepsilon \to 0$. In Section 4 we give the details of the proof.

In Section 5 we tackle the problem of uniqueness of minimizers in the Heisenberg group of arbitrary dimension (see Theorem B). We also generalize the comparison principle in [3] (cf. Theorem C, Theorem C$'$ there) to a weak version and for a wide class of $\vec{F}$'s (see Theorem C below).

Theorem B. *Let Ω be a bounded domain in R^{2n}. Let $u, v \in W^{1,2}(\Omega)$ be two minimizers for $\mathcal{F}(\cdot)$ such that $u - v \in W_0^{1,2}(\Omega)$. Suppose $H \in L^\infty(\Omega)$* and *$\vec{F} \in W^{1,2}(\Omega)$ satisfying $div\vec{F}^* > 0$ (a.e.). Then $u \equiv v$ in Ω (a.e.).*

We remark that in the specific case $\vec{F} = -\vec{X}^*$ the assumptions in Theorem B are satisfied. On the other hand, the condition $div\vec{F}^* > 0$ is essential in Theorem B. Let $\Omega = B_2 - \bar{B}_1 \subset R^2$ where B_r denotes the open ball of radius r. Consider the case $\vec{F} = 0$ and $H = \frac{1}{r}$. Let $u = f(r), v = g(r)$, and $f \neq g$ with the properties that $f(1) = g(1)$, $f(2) = g(2)$, and $f' > 0, g' > 0$ for $1 \leq r \leq 2$. Then it is easy to see that u and v are two minimizers for the associated $\mathcal{F}(\cdot)$ (see also page 162 in [3]).

Theorem C. *Let Ω be a bounded domain in R^{2n}. Let $\vec{F}$ (a vector field) $\in W^{1,2}(\Omega)$ satisfy $div\vec{F}^* > 0$ (a.e.). Suppose $u, v \in W^{1,2}(\Omega)$ satisfy the following conditions:*

$$\begin{aligned} divN(u) &\geq divN(v) \text{ in } \Omega \text{ (in the weak sense);} \\ u &\leq v \text{ on } \partial\Omega. \end{aligned}$$

Then $u \leq v$ *in* Ω.

In Sections 6, we study the relation between C^2-smooth solutions and minimizers. In [3] (Theorem B there), we proved that if u is a C^2-smooth solution to (1.1) in dimension 2 with H bounded near a singular point p_0, then either p_0 is isolated in $S(u)$ or there exists a small neighborhood B of p_0 which intersects with $S(u)$ in exactly a C^1-smooth curve Γ through p_0 (the condition on H can be weaker). Moreover, Γ divides B into two disjoint nonsingular domains B^+ and B^-, and $N(u)(p_0^+) \equiv \lim_{p \in B^+ \to p_0} N(u)(p)$ and $N(u)(p_0^-) \equiv \lim_{p \in B^- \to p_0} N(u)(p)$ exist. Also $N(u)(p_0^+) = -N(u)(p_0^-)$ (see Proposition 3.5 in [3]). In Section 6 and the first part of Section 7 (see Proposition 6.2, Theorem 6.3, (7.1), and (7.2)), we will generalize such a situation and give a criterion for u to be a minimizer. In particular, suppose u is C^2-smooth. Then Proposition 6.2 or Theorem 6.3 gives a criterion for u to be a minimizer in the situation $H_{m-1}(S(u)) > 0$ while if $H_{m-1}(S(u)) = 0$, u must be a minimizer (see Lemma 6.1).

In [16], Pauls constructed two different C^2 (in fact C^∞) smooth solutions to the p-minimal surface equation ((1.1) with $H = 0$ or (1.6) with $H = 0$ and $\vec{F} = -\vec{X}^*$) with the same C^∞-smooth boundary value and the same p-area in $\Omega \subset R^2$. These two solutions do not satisfy the criterion in Proposition 6.2 or Theorem 6.3, hence none of them is a minimizer. We can also see this fact according to Theorem B (uniqueness of minimizers). In Section 7 we construct the actual minimizer for Pauls' example (see Example 7.3).

In dimensions higher than 2, the situation is quite different. The size of the singular set can be relatively small under a suitable condition on $\vec{F} = (F_I)$. For $x \geq 0$, let $[x]$ denote the largest integer less or equal than x. In Section 6 we obtain an estimate for the size of the singular set and a condition on $\vec{F}$ for a C^2-smooth solution to (1.6) to be a minimizer (see Theorems D and E below). Recall that dim_E denotes the Hausdorff dimension with respect to the Euclidean metric.

Theorem D. *Let* Ω *be a domain in* R^m. *Suppose* $u \in C^2(\Omega)$ *and* $F_I \in C^1(\Omega)$. *Then for any* $p \in \Omega$, *there exists a neighborhood* V *of* p *in* Ω *such that* $S(u) \cap V$ *is a submanifold of* V *satisfying*

$$dim_E(S(u) \cap V) \leq m - [\frac{rank\ (\partial_J F_I - \partial_I F_J)(p) + 1}{2}]. \tag{1.9}$$

Theorem E. *Let* Ω *be a bounded domain in* R^m, $m \geq 2$. *Suppose* $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$ *is a* C^2*-smooth solution to (1.6) with* $H \in C^0(\Omega \backslash S(u)) \cap L^\infty(\Omega)$ *and* $F_I \in C^1(\Omega)$. *Suppose there holds*

$$[\frac{rank\ (\partial_J F_I - \partial_I F_J) + 1}{2}] \geq 2 \tag{1.10}$$

for all $p \in \Omega$. *Then* u *is a weak solution to (1.6) and a minimizer for (1.7) if in addition* $u \in W^{1,1}(\Omega)$.

Corollary F. *Let Ω be a bounded domain in R^{2n}. Suppose $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$ is a C^2-smooth solution to the p-minimal surface equation ((1.1) with $H = 0$). Then in dimension ≥ 4 ($n \geq 2$), u is a weak solution to the p-minimal surface equation and a minimizer for (1.2) with $H = 0$ if in addition $u \in W^{1,1}(\Omega)$.*

In Section 8 we study the uniqueness of solutions to elliptic approximating equations $Q_\varepsilon u = H$ (see (4.1)), $\varepsilon > 0$. Since this is an elliptic equation for a given $\varepsilon > 0$, the uniqueness of solutions follows essentially from the known elliptic theory (see e.g. [9]). But for the reader's convenience, we include a proof here as the Appendix.

We were aware of the paper [17] while this work was being done. After this paper was submitted, we were informed of the work [18]. Some problems related to this paper were studied in [17] and [18]. We are grateful to Andrea Malchiodi for many discussions, in particular, in the study of Example 7.3. We would also like to thank the referee for stimulating comments and pointing out many grammatical errors.

2. Hypersurfaces in the Heisenberg group

In this section we introduce some basic notions for a hypersurface in a pseudohermitian manifold. By viewing the Heisenberg group or R^{2n+1} as a suitable pseudohermitian manifold, we give geometric interpretations of (1.1) and (1.4).

Let (M, J, Θ) be a $(2n+1)$-dimensional pseudohermitian manifold with an integrable CR structure J and a global contact form Θ such that the bilinear form $G \equiv \frac{1}{2}d\Theta(\cdot, J\cdot)$ is positive definite on the contact bundle $\xi \equiv \ker \Theta$ ([13]). The metric G is usually called the Levi metric. Consider a hypersurface $\Sigma \subset M$. A point $p \in \Sigma$ is called singular if ξ coincides with $T\Sigma$ at p. Otherwise, p is called nonsingular and $\mathcal{V} \equiv \xi \cap T\Sigma$ is $2n - 1$ dimensional in this case. There is a unique (up to sign) unit vector $N \in \xi$ that is perpendicular to $\mathcal{V}$ with respect to the Levi metric G. We call N the Legendrian normal or the p-normal ("p" stands for "pseudohermitian"). Suppose that Σ bounds a domain Ω in M. We define the p-area $2n$-form $\mathcal{A}$ by computing the first variation ($\mathcal{A}$ will be computed below for the case of the Heisenberg group), away from the singular set, of the standard volume in the p-normal N :

$$\delta_{fN} \int_\Omega \Theta \wedge (d\Theta)^n = c(n) \int_\Sigma f\mathcal{A} \tag{2.1}$$

where f is a C^∞-smooth function on Σ with compact support away from the singular points, and $c(n) = 2^n n!$ is a normalization constant. The sign of N is determined by requiring that $\mathcal{A}$ is positive with respect to the induced orientation on Σ. So we can talk about the p-area of Σ by integrating $\mathcal{A}$ over Σ (which might not be closed from now on). Then we define the p-mean curvature H of Σ as the first variation of the p-area in the direction of N : (the support of f now is also assumed to be away from the boundary of Σ)

$$\delta_{fN} \int_\Sigma \mathcal{A} = -\int_\Sigma fH\mathcal{A}. \tag{2.2}$$

Consider the Heisenberg group viewed as a (flat) pseudohermitian manifold $(R^{2n+1},\ \Theta_0,\ J_0)$. Here $\Theta_0 \equiv dz + \sum_{j=1}^n (x_j dx_{j'} - x_{j'} dx_j)$ at a point $(\vec{X},\ z) \equiv$

$(x_1, x_{1'}, ..., x_n, x_{n'}, z) \in R^{2n+1}$ and $J_0(\mathring{e}_j) \equiv \mathring{e}_{j'}$, $J_0(\mathring{e}_{j'}) \equiv -\mathring{e}_j$ where

$$\mathring{e}_j \equiv \frac{\partial}{\partial x_j} + x_{j'}\frac{\partial}{\partial z}, \quad \mathring{e}_{j'} \equiv \frac{\partial}{\partial x_{j'}} - x_j\frac{\partial}{\partial z} \tag{2.3}$$

$j = 1, 2, ...n$ span $\xi_0 \equiv \ker \Theta_0$. Let Σ be a graph defined by $z = u(\vec{X})$. Note that $\mathring{e}_j$'s and $\mathring{e}_{j'}$'s form an orthonormal basis with respect to the Levi metric $G_0 = (\sum_{j=1}^n dx_j \wedge dx_{j'})\ (\cdot,\ J_0\cdot)$. Observe that an element $v = \sum_{j=1}^n (a_j\mathring{e}_j + b_{j'}\mathring{e}_{j'}) \in \xi_0 \cap T\Sigma$ satisfies $d(z - u(\vec{X}))\ (v) = 0$. It follows that

$$\sum_{j=1}^{n}[(u_{x_j} - x_{j'})a_j + (u_{x_{j'}} + x_j)b_{j'}] = 0. \tag{2.4}$$

Let $N \equiv -D^{-1}\sum_{j=1}^n[(u_{x_j} - x_{j'})\mathring{e}_j + (u_{x_{j'}} + x_j)\mathring{e}_{j'}]$ where $D \equiv (\sum_{j=1}^n[(u_{x_j} - x_{j'})^2 + (u_{x_{j'}} + x_j)^2])^{1/2}$. It is easy to see that N is perpendicular to $\xi_0 \cap T\Sigma$ by (2.4), that it is of the unit length w.r.t. G_0 and hence N is the p-normal (that the associated $\mathcal{A}$ is positive will be shown below). We can now compute $\Theta_0 \wedge (d\Theta_0)^n$ $= c(n)\ dz \wedge dx_1 \wedge dx_{1'} \wedge ... \wedge dx_n \wedge dx_{n'}$ and

$$\iota_N\{\Theta_0 \wedge (d\Theta_0)^n\} = -c(n)D^{-1}\{(I) + (II) + (III)\} \tag{2.5}$$

where ι_N means taking the interior product with N and ($d\hat{x}_I$ deleted)

$$\begin{aligned}
(I) &= \sum_{j=1}^{n}[(u_{x_j} - x_{j'})x_{j'} - (u_{x_{j'}} + x_j)x_j]dx_1 \wedge dx_{1'} \wedge ... \wedge dx_n \wedge dx_{n'} \\
(II) &= -\sum_{j=1}^{n}(u_{x_j} - x_{j'})dz \wedge dx_1 \wedge dx_{1'}...d\hat{x}_j \wedge dx_{j'} ... \wedge dx_n \wedge dx_{n'} \\
(III) &= \sum_{j=1}^{n}(u_{x_{j'}} + x_j)dz \wedge dx_1 \wedge dx_{1'}...dx_j \wedge d\hat{x}_{j'} ... \wedge dx_n \wedge dx_{n'}.
\end{aligned}$$

It follows that

$$\begin{aligned}
&\delta_{fN}\int_\Omega \Theta_0 \wedge (d\Theta_0)^n \\
&= \int_\Omega L_{fN}\{\Theta_0 \wedge (d\Theta_0)^n\} = \int_\Omega d(\iota_{fN}\{\Theta_0 \wedge (d\Theta_0)^n\}) \\
&= \int_\Sigma f\iota_N\{\Theta_0 \wedge (d\Theta_0)^n\}
\end{aligned} \tag{2.6}$$

by the formula $L_v = \iota_v \circ d + d \circ \iota_v$ and Stokes' theorem. Substituting (2.5) into (2.6) and comparing (2.6) with (2.1) gives

$$\mathcal{A} = -D^{-1}\{(I) + (II) + (III)\} \tag{2.7}$$

which simplifies to $Ddx_1 \wedge dx_{1'} \wedge ... \wedge dx_n \wedge dx_{n'}$ on Σ ($z = u(\vec{X})$). Next we compute

$$\delta_{fN}\int_\Sigma \mathcal{A} = \int_\Sigma L_{fN}\mathcal{A} = \int_\Sigma \iota_{fN} \circ d\mathcal{A}. \tag{2.8}$$

Here we have used Stokes' theorem and the condition that the support of f is away from the singular set and the boundary of Σ. Noting that $D = |\nabla u - \vec{X}^*|$ where $\vec{X}^* = (x_{1'}, -x_1, x_{2'}, -x_2, ..., x_{n'}, -x_n)$, we can easily deduce that $d(D^{-1}(I)) = 0$ and

$$d\{D^{-1}[(II) + (III)]\} = (div\frac{\nabla u - \vec{X}^*}{|\nabla u - \vec{X}^*|})dz \wedge dx_1 \wedge dx_{1'} \wedge ... \wedge dx_n \wedge dx_{n'}.$$

It follows from (2.7) and (2.5) that

$$\iota_N \circ d\mathcal{A} = -(div\frac{\nabla u - \vec{X}^*}{|\nabla u - \vec{X}^*|})\mathcal{A}. \tag{2.9}$$

Substituting (2.9) into (2.8) and comparing (2.8) with (2.2), we obtain the following expression for the p-mean curvature H_Σ of the graph $\Sigma = \{(\vec{X}, u(\vec{X}))\}$:

$$H_\Sigma = div\frac{\nabla u - \vec{X}^*}{|\nabla u - \vec{X}^*|}. \tag{2.10}$$

Next we consider a general vector field $\vec{F} = (F_I)$ instead of $-\vec{X}^*$. Let $\Theta_{\vec{F}} \equiv dz + \sum_I F_I dx_I$ where I ranges over 1, $1'$, ..., n, n'. Assume that $\Theta_{\vec{F}}$ is a contact form, i.e., $\Theta_{\vec{F}} \wedge (d\Theta_{\vec{F}})^n \neq 0$ everywhere (satisfied for $\vec{F} = -\vec{X}^*$ as shown previously). For instance, the condition is equivalent to $\partial F_{1'}/\partial x_1 - \partial F_1/\partial x_{1'} \neq 0$ in the case $n = 1$. Define

$$e_I = \frac{\partial}{\partial x_I} - F_I\frac{\partial}{\partial z}, \quad I = 1, 1', ..., n, n'. \tag{2.11}$$

It is easy to see that $\Theta_{\vec{F}}$ annihilates the e_I's. Define the CR structure $J_{\vec{F}}$ on the contact bundle $\ker\Theta_{\vec{F}}$ by $J_{\vec{F}}(e_j) = e_{j'}$ and $J_{\vec{F}}(e_{j'}) = -e_j$ for $j = 1, 2, ..., n$. For the 2-dimensional case ($n = 1$), we can find a nonvanishing scalar function λ ($= 2(\partial F_{1'}/\partial x_1 - \partial F_1/\partial x_{1'})^{-1}$) such that $\{e_1, e_{1'}\}$ forms an orthonormal basis with respect to the Levi metric $G_{\vec{F}}$ associated to $(J_{\vec{F}}, \lambda\Theta_{\vec{F}})$. Let $\psi \equiv z - u(x_1, x_{1'})$ be a defining function for the graph of u. By a formula in Section 2 of [3], we can compute the p-mean curvature $H_{\vec{F}}$ with respect to the pseudohermitian structure $(J_{\vec{F}}, \lambda\Theta_{\vec{F}})$ as follows:

$$\begin{aligned} H_{\vec{F}} &= -div_b\frac{\nabla_b\psi}{|\nabla_b\psi|_{G_{\vec{F}}}} \\ &= -e_1(\frac{e_1\psi}{D_{\vec{F}}}) - e_{1'}(\frac{e_{1'}\psi}{D_{\vec{F}}}) \\ &= \frac{\partial}{\partial x_1}(\frac{u_{x_1} + F_1}{|\nabla u + \vec{F}|}) + \frac{\partial}{\partial x_{1'}}(\frac{u_{x_{1'}} + F_{1'}}{|\nabla u + \vec{F}|}) \\ &= div\frac{\nabla u + \vec{F}}{|\nabla u + \vec{F}|}. \end{aligned} \tag{2.12}$$

Here we have used $|\nabla_b\psi|_{G_{\vec{F}}} = \sqrt{(e_1\psi)^2 + (e_{1'}\psi)^2} = |\nabla u + \vec{F}|$ by (2.11).

3. Minimizers in the Heisenberg group

In this section we deduce some properties of a minimizer in the Heisenberg group. In fact we consider a more general area functional (this is just (1.7)):

$$\mathcal{F}(u) \equiv \int_\Omega \{\mid \nabla u + \vec{F} \mid + Hu\} \tag{3.1}$$

where $\Omega \subset R^m$ is a bounded domain, $\vec{F}$ is an arbitrary (say, L^1) vector field on Ω, and $H \in L^\infty(\Omega)$ (we omit the Euclidean volume element).

Definition 3.1. $u \in W^{1,1}(\Omega)$ is called a minimizer for $\mathcal{F}(u) \equiv \int_\Omega \{|\nabla u + \vec{F}| + Hu\}$ if $\mathcal{F}(u) \leq \mathcal{F}(u+\varphi)$ for any $\varphi \in W_0^{1,1}(\Omega)$, where $\vec{F} \in L^1(\Omega)$ and $H \in L^\infty(\Omega)$.

We are going to investigate the first variation of $\mathcal{F}$. Let $u, \varphi \in W^{1,1}(\Omega)$ and $u_\varepsilon \equiv u + \varepsilon\varphi$ for $\varepsilon \in R$. It follows that $u_\varepsilon - u_{\hat\varepsilon} = (\varepsilon - \hat\varepsilon)\varphi$. Let $S(u_\varepsilon)$, the singular set of u_ε, denote the set of points where $\nabla u_\varepsilon + \vec{F} = 0$. So from (3.1) (noting that $\mid \nabla u_\varepsilon + \vec{F} \mid = \mid \varepsilon - \hat\varepsilon \mid \mid \nabla\varphi \mid$ on $S(u_{\hat\varepsilon})$) we have

$$\begin{aligned} \mathcal{F}(u_\varepsilon) \quad = \quad & \mid \varepsilon - \hat\varepsilon \mid \int_{S(u_{\hat\varepsilon})} \mid \nabla\varphi \mid + \int_{\Omega\setminus S(u_{\hat\varepsilon})} \mid \nabla u_\varepsilon + \vec{F} \mid \\ & + \int_\Omega H u_{\hat\varepsilon} + \int_\Omega (\varepsilon - \hat\varepsilon) H\varphi. \end{aligned} \tag{3.2}$$

Since $\mid \nabla u_\varepsilon + \vec{F} \mid^2 - \mid \nabla u_{\hat\varepsilon} + \vec{F} \mid^2 = 2(\varepsilon - \hat\varepsilon)(\nabla u_{\hat\varepsilon} + \vec{F}) \cdot \nabla\varphi + (\varepsilon - \hat\varepsilon)^2 \mid \nabla\varphi \mid^2$, we compute from (3.2)

$$\begin{aligned} \frac{\mathcal{F}(u_\varepsilon) - \mathcal{F}(u_{\hat\varepsilon})}{\varepsilon - \hat\varepsilon} \quad = \quad & \frac{|\varepsilon - \hat\varepsilon|}{\varepsilon - \hat\varepsilon} \int_{S(u_{\hat\varepsilon})} |\nabla\varphi| + \int_{\Omega\setminus S(u_{\hat\varepsilon})} \frac{2(\nabla u_{\hat\varepsilon} + \vec{F}) \cdot \nabla\varphi + (\varepsilon - \hat\varepsilon)|\nabla\varphi|^2}{|\nabla u_\varepsilon + \vec{F}| + |\nabla u_{\hat\varepsilon} + \vec{F}|} \\ & + \int_\Omega H\varphi. \end{aligned}$$

Note that the integrand of the middle term in the right-hand side of the above formula actually equals $(\varepsilon - \hat\varepsilon)^{-1}$ $(\mid \nabla u_\varepsilon + \vec{F} \mid - \mid \nabla u_{\hat\varepsilon} + \vec{F} \mid)$ whose absolute value is less than or equal to $\mid \nabla\varphi \mid$. Therefore by Lebesque's dominated convergence theorem, we can easily take the limit as $\varepsilon \to \hat\varepsilon\pm$ (+: the right-hand limit; −: the left-hand limit), and obtain

$$\frac{d\mathcal{F}(u_{\hat\varepsilon\pm})}{d\varepsilon} = \pm \int_{S(u_{\hat\varepsilon})} \mid \nabla\varphi \mid + \int_{\Omega\setminus S(u_{\hat\varepsilon})} N(u_{\hat\varepsilon}) \cdot \nabla\varphi + \int_\Omega H\varphi \tag{3.3}$$

where $N(v) \equiv \frac{\nabla v + \vec{F}}{|\nabla v + \vec{F}|}$ is defined on $\Omega\setminus S(v)$. Note that $N(u_{\hat\varepsilon}) \cdot \nabla\varphi \in L^1(\Omega\setminus S(u_{\hat\varepsilon}))$ since $|N(u_{\hat\varepsilon}) \cdot \nabla\varphi| \leq |N(u_{\hat\varepsilon})|\ |\nabla\varphi| = |\nabla\varphi|$ and $\nabla\varphi \in L^1(\Omega)$ by the assumption. Also from the above argument, we have the estimate

$$\frac{\mid \mathcal{F}(u_\varepsilon) - \mathcal{F}(u_{\hat\varepsilon}) \mid}{\mid \varepsilon - \hat\varepsilon \mid} \leq \int_\Omega \mid \nabla\varphi \mid + ||H||_\infty \int_\Omega |\varphi|.$$

Namely, $\mathcal{F}(u_\varepsilon)$ is Lipschitz continuous in ε for $\varphi \in W^{1,1}(\Omega)$. Let $\kappa(\varepsilon)$ denote the Lebesque measure of the set $S(u_\varepsilon) \cap \{\nabla\varphi \neq 0\}$. We claim that there are at most countably many ε's with $\kappa(\varepsilon) > 0$ for a fixed φ. First observe that $S(u_{\varepsilon_1}) \cap S(u_{\varepsilon_2}) \subset \{\nabla\varphi = 0\}$, and hence $(S(u_{\varepsilon_1}) \cap \{\nabla\varphi \neq 0\}) \cap (S(u_{\varepsilon_2}) \cap \{\nabla\varphi \neq 0\}) = \emptyset$ (empty). Let $|\Omega|$ denote the volume of the bounded domain Ω. So the number of ε such that $\kappa(\varepsilon) > \frac{1}{n}$ for any positive integer is at most $[n|\Omega|] + 1$ where $[x]$ denotes the largest integer less than or equal to x. Therefore there are at most countably many ε's with $\kappa(\varepsilon) > 0$. We call such an ε singular, otherwise regular (i.e., $\kappa(\varepsilon) = 0$). By (3.3), we obtain (3.4) in the following Lemma.

Lemma 3.1. *(1) $\mathcal{F}(u_\varepsilon)$ is Lipschitz continuous in ε for $\varphi \in W^{1,1}(\Omega)$. (2) There are at most countably many singular ε's. (3) For a regular ε, $\int_{S(u_\varepsilon)} \mid \nabla\varphi \mid= 0$, $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon}$ exists, and*

$$\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} = \int_{\Omega\setminus S(u_\varepsilon)} N(u_\varepsilon)\cdot\nabla\varphi + \int_\Omega H\varphi. \tag{3.4}$$

Next for $\varepsilon_2, \varepsilon_1$ regular with $\varepsilon_2 > \varepsilon_1$, we compute the difference of $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon}$ for $\varepsilon = \varepsilon_2, \varepsilon_1$ by (3.4). Using $\kappa(\varepsilon_j) = 0$, $j = 1, 2$ to shrink the domain of the integral, we obtain

$$\frac{d\mathcal{F}(u_{\varepsilon_2})}{d\varepsilon} - \frac{d\mathcal{F}(u_{\varepsilon_1})}{d\varepsilon} = \int_{\Omega\setminus[S(u_{\varepsilon_2})\cup S(u_{\varepsilon_1})]} [N(u_{\varepsilon_2}) - N(u_{\varepsilon_1})]\cdot\nabla\varphi \geq 0. \tag{3.5}$$

Here we have used Lemma 5.1′ (also holds for $u, v \in W^{1,1}$) in [3] to conclude the last inequality in (3.5) by noting that $\nabla\varphi = (\varepsilon_2 - \varepsilon_1)^{-1}(\nabla u_{\varepsilon_2} - \nabla u_{\varepsilon_1})$. We have the following result.

Lemma 3.2. *(1) $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon}$ is an increasing function of ε for ε regular. (2) Let ε_j, $j = 1, 2, ...$, be a sequence of decreasing (increasing, respectively) regular numbers tending to $\hat{\varepsilon}$ ($\hat{\varepsilon}$ may be singular) as $j \to \infty$. Then we have*

$$\lim_{j\to\infty} \frac{d\mathcal{F}(u_{\varepsilon_j})}{d\varepsilon} = \frac{d\mathcal{F}(u_{\hat{\varepsilon}+})}{d\varepsilon} \quad \left(= \frac{d\mathcal{F}(u_{\hat{\varepsilon}-})}{d\varepsilon}, \text{ respectively}\right). \tag{3.6}$$

Note that we have the precise expressions for the right-hand limit $\frac{d\mathcal{F}(u_{\hat{\varepsilon}+})}{d\varepsilon}$ and the left-hand limit $\frac{d\mathcal{F}(u_{\hat{\varepsilon}-})}{d\varepsilon}$ at $\hat{\varepsilon}$ in (3.3).

Proof. (1) follows from (3.5). To prove (2), first observe that $\int_{S(u_{\varepsilon_j})} |\nabla\varphi| = 0$ by the definition of ε_j being regular. Therefore we have

$$\int_{\cup_{j=1}^\infty S(u_{\varepsilon_j})} |\nabla\varphi| = 0. \tag{3.7}$$

Let $S_\infty \equiv \cup_{j=1}^\infty S(u_{\varepsilon_j})$. Since $|N(u_{\varepsilon_j})| \leq 1$, we estimate $\mid \int_{S_\infty} N(u_{\varepsilon_j}) \cdot \nabla\varphi \mid \leq \int_{S_\infty} |\nabla\varphi| = 0$ by (3.7). So we obtain

$$\int_{S_\infty} N(u_{\varepsilon_j}) \cdot \nabla\varphi = 0. \tag{3.8}$$

It then follows from (3.4) and (3.8) that

$$\begin{aligned}\frac{d\mathcal{F}(u_{\varepsilon_j})}{d\varepsilon} &= \int_{\Omega\backslash S(u_{\varepsilon_j})} N(u_{\varepsilon_j})\cdot\nabla\varphi+\int_\Omega H\varphi \\ &= \int_{\Omega\backslash S_\infty} N(u_{\varepsilon_j})\cdot\nabla\varphi+\int_\Omega H\varphi.\end{aligned}\tag{3.9}$$

On the other hand, observe that $\lim_{j\to\infty} N(u_{\varepsilon_j}) = N(u_{\hat\varepsilon})$ in $\Omega\backslash[S_\infty\cup S(u_{\hat\varepsilon})]$ and

$$N(u_{\varepsilon_j}) = \frac{(\nabla u_{\hat\varepsilon}+\vec F)+(\varepsilon_j-\hat\varepsilon)\nabla\varphi}{|(\nabla u_{\hat\varepsilon}+\vec F)+(\varepsilon_j-\hat\varepsilon)\nabla\varphi|} = \frac{(\varepsilon_j-\hat\varepsilon)\nabla\varphi}{|\varepsilon_j-\hat\varepsilon||\nabla\varphi|}\tag{3.10}$$

in $S(u_{\hat\varepsilon})\backslash S_\infty$. Now we compute

$$\begin{aligned}&\int_{\Omega\backslash S_\infty} N(u_{\varepsilon_j})\cdot\nabla\varphi \\ =\ &(\int_{S(u_{\hat\varepsilon})\backslash S_\infty}+\int_{\Omega\backslash[S_\infty\cup S(u_{\hat\varepsilon})]})N(u_{\varepsilon_j})\cdot\nabla\varphi \\ =\ &\frac{\varepsilon_j-\hat\varepsilon}{|\varepsilon_j-\hat\varepsilon|}\int_{S(u_{\hat\varepsilon})\backslash S_\infty}|\nabla\varphi|+\int_{\Omega\backslash[S_\infty\cup S(u_{\hat\varepsilon})]} N(u_{\varepsilon_j})\cdot\nabla\varphi \\ \to\ &\pm\int_{S(u_{\hat\varepsilon})}|\nabla\varphi|+\int_{\Omega\backslash S(u_{\hat\varepsilon})} N(u_{\hat\varepsilon})\cdot\nabla\varphi\end{aligned}\tag{3.11}$$

as $j\to\infty$ (+ for decreasing ε_j; − for increasing ε_j). Here we have used (3.10) and Lebesque's dominated convergence theorem. By (3.9), (3.11), and in view of (3.3), we have proved (3.6).

Q.E.D.

Definition 3.2. Let $\Omega\subset R^m$ be a bounded domain. Let $\vec F$ be an L^1_{loc} vector field on Ω. Let $H\in L^1_{loc}(\Omega)$. We say $u\in W^1(\Omega)$ is a weak solution to the equation (1.6), i.e., $divN(u) = H$ in Ω if and only if for any $\varphi\in C_0^\infty(\Omega)$, there holds

$$\int_{S(u)}|\nabla\varphi|+\int_{\Omega\backslash S(u)} N(u)\cdot\nabla\varphi+\int_\Omega H\varphi\geq 0.\tag{3.12}$$

Recall that $N(u)\equiv\frac{\nabla u+\vec F}{|\nabla u+\vec F|}$, $S(u)\equiv\{\nabla u+\vec F=0\}$, and $N(u)\cdot\nabla\varphi\in L^1(\Omega\backslash S(u))$ since $|N(u)\cdot\nabla\varphi|\leq|N(u)|\,|\nabla\varphi| = |\nabla\varphi|$ and $\nabla\varphi\in L^1(\Omega)$ by assumption.. Note that with φ replaced by $-\varphi$ in (3.12), we also have $-\int_{S(u)}|\nabla\varphi| + \int_{\Omega\backslash S(u)} N(u)\cdot\nabla\varphi$ $+\int_\Omega H\varphi\leq 0$. Moreover, if the $(m-1)$-dimensional Hausdorff measure of $S(u)$ vanishes, then the equality holds in (3.12). We remark that in Definition 3.2 for the case $H\in L^\infty(\Omega)$, the space $C_0^\infty(\Omega)$ of test functions can be replaced by $W_0^{1,1}(\Omega)$ since the former is dense in the latter in the $W^{1,1}$ norm ([9]). Note that in the definition of a minimizer, we require $u\in W^{1,1}(\Omega)$, $\vec F\in L^1(\Omega)$, and $H\in L^\infty(\Omega)$ while for the definition of a weak solution, u can be in a larger space $W^1(\Omega)$, $\vec F\in L^1_{loc}(\Omega)$, and $H\in L^1_{loc}(\Omega)$.

Theorem 3.3. *Let $u \in W^{1,1}(\Omega)$, $\vec{F} \in L^1(\Omega)$, and $H \in L^\infty(\Omega)$. Then u is a minimizer for $\mathcal{F}(\cdot)$ if and only if u is a weak solution to the equation $divN(u) = H$.*

Proof. Suppose u is a minimizer for $\mathcal{F}(u)$. Then $\frac{d\mathcal{F}(u_{0+})}{d\varepsilon} \geq 0$, and hence (3.12) follows from (3.3) (letting $\hat{\varepsilon} = 0$ in (3.3)). So u is a weak solution. Conversely, suppose u is a weak solution. Since $\mathcal{F}(u_\varepsilon)$ is Lipschitz continuous in ε by Lemma 3.1 (1), $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon}$ exists a.e. (in fact at least for regular ε) and it is integrable. Moreover, we have

$$\mathcal{F}(u+\varphi) - \mathcal{F}(u) = \int_0^1 \frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} d\varepsilon. \tag{3.13}$$

On the other hand, from Lemma 3.2 and the definition of weak solution (Definition 3.2), we obtain that $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} \geq 0$ for any regular $\varepsilon \in [0,1]$ in view of (3.3) (take $\hat{\varepsilon} = 0$). By Lemma 3.1 (2), $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} \geq 0$ a.e.. It follows from (3.13) that $\mathcal{F}(u+\varphi) \geq \mathcal{F}(u)$. That is to say, u is a minimizer for $\mathcal{F}(u)$.

Q.E.D.

4. EXISTENCE OF MINIMIZERS-PROOF OF THEOREM A

Let Ω be a bounded domain in $R^m, m \geq 2$. Consider the following elliptic approximation $u = u_\varepsilon$ ($\varepsilon > 0$) (a geometric interpretation can be found in [16]) with given boundary value φ ($\in C^{2,\alpha}(\bar{\Omega})$, $0 < \alpha < 1$, say) :

$$\begin{aligned} Q_\varepsilon u &\equiv div\left(\frac{\nabla u + \vec{F}}{\sqrt{\varepsilon^2 + |\nabla u + \vec{F}|^2}}\right) = 0 \quad \text{in } \Omega, \\ u &= \varphi \quad \text{on } \partial\Omega \end{aligned} \tag{4.1}$$

where $\vec{F} = (F_I)$, $I = 1, ..., m$. In the case of $m = 2n$, I ranges over 1, $1'$, ..., n, n' (e.g., $F_I = -x_{I'}$ for the case of a p-minimal surface. Here we use the convention that $x_{j''} = -x_j$, $j = 1$, ..., n). We will make use of Theorem 11.8 in [9] to solve (4.1) in $C^{2,\alpha}(\bar{\Omega})$ (then a subsequence of u_ε will converge to what we want). First we check that Q_ε is elliptic. A direct computation shows that (summation convention applies)

$$\begin{aligned} Q_\varepsilon u &= \frac{u_{II}(\varepsilon^2 + |\nabla u + \vec{F}|^2) - (u_I + F_I)(u_J + F_J)u_{IJ}}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}} \\ &\quad + \frac{(\varepsilon^2 + |\nabla u + \vec{F}|^2)\partial_I F_I - (u_I + F_I)(u_J + F_J)\partial_I F_J}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}} \\ &= a_{IJ}(\varepsilon, x, \nabla u)u_{IJ} + b(\varepsilon, x, \nabla u) \end{aligned} \tag{4.2}$$

where

$$a_{IJ}(\varepsilon, x, \nabla u) = \frac{\delta_{IJ}(\varepsilon^2 + |\nabla u + \vec{F}|^2) - (u_I + F_I)(u_J + F_J)}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}} \tag{4.3}$$

and

$$b(\varepsilon, x, \nabla u) = \frac{(\varepsilon^2 + |\nabla u + \vec{F}|^2)\partial_I F_I - (u_I + F_I)(u_J + F_J)\partial_I F_J}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}}.$$

For $0 \neq (p_I) \in R^m$, we compute from (4.3) that

$$\begin{aligned} a_{IJ}p_I p_J &= \frac{(\varepsilon^2 + |\nabla u + \vec{F}|^2)p_I^2 - (u_I + F_I)(u_J + F_J)p_I p_J}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}} \\ &\geq \frac{\varepsilon^2 p_I^2}{[\varepsilon^2 + |\nabla u + \vec{F}|^2]^{3/2}} > 0. \end{aligned} \tag{4.4}$$

Here we have used Cauchy's inequality $|(\nabla u + \vec{F}) \cdot (p_I)|^2 \leq |\nabla u + \vec{F}|^2 p_I^2$ (noting that p_I^2 means the sum $\Sigma_I p_I^2$). It follows from (4.2) and (4.4) that Q_ε is elliptic.

To apply Theorem 11.8 in [9], we need to get an apriori estimate in $C^1(\bar{\Omega})$ -norm at least. Suppose u_ε is a $C^{2,\alpha}(\bar{\Omega})$ solution to the equation $Q_\epsilon u = 0$ (assuming $\vec{F} \in C^{1,\alpha}(\bar{\Omega})$; later replacing $\vec{F}$ by $\sigma\vec{F}$), $u = \sigma\varphi$ on $\partial\Omega$, $0 \leq \sigma \leq 1$. In the case of $F_I = -x_{I'}$, $\partial_I F_I = 0$, $(u_I + F_I)(u_J + F_J)\partial_I F_J = -(u_{J'} + F_{J'})(u_J + F_J) = 0$, and hence $b(\varepsilon, x, \nabla u) = 0$. Since Q_ε is elliptic, it follows from the maximum principle (see e.g. Problem 10.1 in [9]) that

$$\sup_{\Omega} |u_\varepsilon| \leq \sup_{\partial\Omega} |u_\varepsilon| = \sup_{\partial\Omega} |\sigma\varphi| \leq \sup_{\partial\Omega} |\varphi|. \tag{4.5}$$

Note that the right hand side is independent of ε. For a general $\vec{F}$, we will invoke the comparison principle for a second order, quasilinear operator with a "tail" term (namely, Theorem 10.1 in [9]). First we can find the comparison functions as shown below. Let $|| \ ||_\infty$ denote the supremum norm. Let B_R denote the ball of radius R, centered at the origin.

Lemma 4.1. *Let $\Omega \subset B_R \subset R^m$ be a bounded domain. Suppose $\vec{F} \in C^1(\Omega)$ be such that F_I and $\partial_I F_J$ are all bounded in Ω. Then there are C^∞-smooth functions*

$$w = e^{x_1 + \kappa R} + e^{x_2 + \kappa R}, \ w' = -e^{x_1 + \kappa' R} - e^{x_2 + \kappa' R}$$

in R^m, where $\kappa = \kappa(\varepsilon, R, ||F_I||_\infty, ||\partial_I F_J||_\infty) > 0$ and $\kappa' = \kappa'(\varepsilon, R, ||F_I||_\infty, ||\partial_I F_J||_\infty) > 0$, such that $Q_\varepsilon w > 0$ and $Q_\varepsilon w' < 0$ in Ω. Moreover, we can choose κ and κ' independent of ε (but depending on ε_0) for $0 < \varepsilon \leq \varepsilon_0$, a positive constant.

Proof. Let w have the above expression with κ to be determined later. Let $w_1 \equiv \partial_{x_1} w$, $w_{11} \equiv \partial_{x_1}^2 w$, $w_{12} \equiv \partial_{x_2}\partial_{x_1} w$, and so on. It follows that

$$\begin{aligned} w_{11} &= w_1 = e^{x_1 + \kappa R}, \ w_{22} = w_2 = e^{x_2 + \kappa R}, \text{ and} \\ w_{IJ} &= 0, \text{ otherwise.} \end{aligned} \tag{4.6}$$

In view of (4.2) with u replaced by w, we compute the dominating (will be clear soon) term in the numerator, which is cubic in w as follows:

$$
\begin{aligned}
(4.7)\qquad & (\sum_J w_J^2)(\sum_I w_{II}) - \sum_{I,J} w_I w_J w_{IJ} \\
= \; & w_1^2 w_{22} + w_2^2 w_{11} \quad \text{(by (4.6))} \\
= \; & e^{2x_1+x_2+3\kappa R} + e^{2x_2+x_1+3\kappa R} \quad \text{(by (4.6))}.
\end{aligned}
$$

It is easy to see that any other term in the expansion of the numerator is bounded by either $c_1 e^{2\kappa R}$, $c_2 e^{\kappa R}$ or c_3 for κ large. Here $c_i = c_i(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty)$, $i = 1,\ 2,\ 3$, are independent of κ. Therefore we have $Q_\varepsilon w > 0$ in Ω by (4.7) for a large $\kappa = \kappa(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty)$. Moreover, κ is independent of ε for $0 < \varepsilon \leq \varepsilon_0$, a positive constant. Similarly, we can find $\kappa' = \kappa'(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty) > 0$ such that $Q_\varepsilon w' < 0$ in Ω.

Q.E.D.

Proposition 4.2. *Let $\Omega \subset B_R \subset R^m$ be a bounded domain. Let $\vec{F} \in C^1(\Omega)$ such that F_I and $\partial_I F_J$ are all bounded in Ω. Suppose $u_\varepsilon \in C^2(\Omega) \cap C^0(\bar{\Omega})$ satisfies (4.1), i.e., $Q_\varepsilon u_\varepsilon = 0$ in Ω and $u_\varepsilon = \sigma\varphi \in C^0$, $0 \leq \sigma \leq 1$, on $\partial\Omega$. Then there exists a constant $C = C(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty,\ ||\varphi||_\infty)$ (independent of σ) such that*

$$
\sup_\Omega |u_\varepsilon| \leq C. \tag{4.8}
$$

Moreover, the bounds hold uniformly for $0 < \varepsilon \leq \varepsilon_0$, a positive constant.

Proof. Let w, w' be the comparison functions as in Lemma 4.1. On $\partial\Omega$, $w \leq \sigma\varphi + C_1 = u_\varepsilon + C_1$ for some constant $C_1 = C_1(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty,\ ||\varphi||_\infty)$ (independent of σ) independent of ε for $0 < \varepsilon \leq \varepsilon_0$, a positive constant. On the other hand, we have $Q_\varepsilon w > 0 = Q_\varepsilon(u_\varepsilon + C_1)$ in Ω by Lemma 4.1 and the observation that $Q_\varepsilon(u_\varepsilon + C_1) = Q_\varepsilon u_\varepsilon$. Now we apply the comparison principle for quasilinear operators (e.g. Theorem 10.1 in [9]) to conclude that

$$
w \leq u_\varepsilon + C_1 \text{ in } \Omega. \tag{4.9}
$$

Similarly, there is a constant $C_2 = C_2(\varepsilon,\ R,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty,\ ||\varphi||_\infty)$ (independent of σ) independent of ε for $0 < \varepsilon \leq \varepsilon_0$, such that $w' \geq \sigma\varphi - C_2 = u_\varepsilon - C_2$ on $\partial\Omega$ and $Q_\varepsilon w' < 0 = Q_\varepsilon(u_\varepsilon - C_2)$ in Ω. So we obtain from the comparison principle that

$$
w' \geq u_\varepsilon - C_2 \text{ in } \Omega. \tag{4.10}
$$

Thus (4.8) follows from (4.9) and (4.10).

Q.E.D.

For the gradient estimate, we will reduce the problem to a gradient estimate at the boundary. We need to require a condition on $\vec{F}$. Suppose there are C^1-smooth functions f_K's ($K = 1, ..., m$) in Ω such that

$$
\partial_K F_I = \partial_I f_K. \tag{4.11}
$$

We remark that if both F_I and G_I satisfy the condition (4.11), so does $F_I + G_I$. In fact, we can write down all the (local) solutions to (4.11). It is easy to see from (4.11) that $\partial_K\ (\partial_J F_I - \partial_I F_J) = 0$ for all I, J, $K = 1, ..., m$. It follows that

$$\partial_J F_I - \partial_I F_J = C_{IJ} \tag{4.12}$$

where the constants C_{IJ} satisfy the skew- symmetric relation: $C_{IJ} = -C_{JI}$. Since the left-hand side of (4.12) is linear in $\vec{F}$, the general solutions are the solutions to $\partial_J F_I - \partial_I F_J = 0$ plus a special solution. Let $\omega \equiv \sum_I F_I dx_I$. Then $d\omega = 0$ if $\partial_J F_I - \partial_I F_J = 0$. So locally there is a function g such that $\omega \equiv dg$. Hence $F_I = \partial_I g$. On the other hand, we observe that $\tilde{F_I} \equiv \frac{1}{2}\sum_K C_{IK} x_K$ is a special solution to (4.12). So the general solutions to (4.12) are

$$F_I = \partial_I g + \frac{1}{2}\sum_K C_{IK} x_K. \tag{4.13}$$

It is then easy to verify that $\vec{F} = (F_I)$ having the form (4.13) are also solutions to (4.11) for $f_K = \partial_K g + \frac{1}{2}\sum_J C_{JK} x_J$.

Proposition 4.3. *Let $\Omega \subset R^m$ be a bounded domain. Let $\vec{F} = (F_I) \in C^1(\Omega)$ satisfy the condition (4.11), for all $I, K = 1, ..., m$, where all f_K's are bounded. Suppose $u_\varepsilon \in C^2(\bar{\Omega})$ satisfies the equation $Q_\varepsilon u_\varepsilon = H_0$, a constant, in Ω. Then we have*

$$\sup_{\Omega} |\partial_K u_\varepsilon| \leq \sup_{\partial\Omega} |\partial_K u_\varepsilon| + 2||f_K||_\infty. \tag{4.14}$$

Proof. Write $\nabla u + \vec{F} = (u_I + F_I)$. Let $D_\varepsilon(u) \equiv \sqrt{\varepsilon^2 + |\nabla u + \vec{F}|^2}$. Compute (summing over J while fixing I and K)

$$\begin{aligned} &\partial_K \frac{u_I + F_I}{D_\varepsilon(u)} \\ = \quad &\frac{u_{IK} + \partial_K F_I}{D_\varepsilon(u)} - \frac{(u_I + F_I)(u_J + F_J)(u_{JK} + \partial_K F_J)}{D_\varepsilon^3(u)} \\ = \quad &\frac{\delta_{IJ} - \nu_I(u)\nu_J(u)}{D_\varepsilon(u)} \partial_J (u_K + f_K) \end{aligned} \tag{4.15}$$

where $\nu_I(u) \equiv (u_I + F_I)/D_\varepsilon(u)$ and we have used the condition (4.11). Now for $v \in C_0^2(\Omega)$, we compute

$$
\begin{aligned}
(4.16)\,0 &= \int_\Omega (Q_\varepsilon u_\varepsilon - H_0)\partial_K v = \int_\Omega \partial_I \frac{(u_\varepsilon)_I + F_I}{D_\varepsilon(u_\varepsilon)}\partial_K v \text{ (summing over } I)\\
&= -\int_\Omega \frac{(u_\varepsilon)_I + F_I}{D_\varepsilon(u_\varepsilon)}\partial_I\partial_K v\\
&= \int_\Omega \partial_K \frac{(u_\varepsilon)_I + F_I}{D_\varepsilon(u_\varepsilon)}\partial_I v \quad (\partial_I\partial_K = \partial_K\partial_I)\\
&= \int_\Omega \{a_{IJ}(\varepsilon, x, \nabla u_\varepsilon)\partial_J[(u_\varepsilon)_K + f_K]\}\partial_I v \text{ (summing over } I \text{ and } J)
\end{aligned}
$$

by (4.15) with u replaced by u_ε. Here $a_{IJ}(\varepsilon, x, \nabla u_\varepsilon) = [\delta_{IJ} - \nu_I(u_\varepsilon)\nu_J(u_\varepsilon)]/D_\varepsilon(u_\varepsilon)$ (cf. (4.3)). It is then easy to see that (4.16) holds also for $v \in C_0^1(\Omega)$ (use the regularization v_h of (7.13) in [9] to approximate v). So $(u_\varepsilon)_K + f_K$ is a weak solution to the equation $Lw \equiv \partial_I\{a_{IJ}(\varepsilon, x, \nabla u_\varepsilon)\partial_J w\} = 0$ (cf. (8.2) in [9]). By (4.4), this is an elliptic equation in divergence form. So by the maximum principle (e.g. Theorem 8.1 in [9] with $b^i = c^i = d = 0$ and a_{IJ} bounded), we have

$$\sup_\Omega |(u_\varepsilon)_K + f_K| \le \sup_{\partial\Omega} |(u_\varepsilon)_K + f_K|.$$

Then (4.14) follows.

Q.E.D.

For a general $\vec{F}$, the bound for ∇u_ε may depend on ε if we invoke the maximum principle for a more general situation (for instance, Theorem 8.16 in [9]).

To perform the boundary gradient estimate, we need a comparison function to apply the comparison principle. Let $\Omega \subset R^m$ be a bounded domain with coordinates denoted by $x_1, x_2, ..., x_m$. We call a coordinate system orthonormal if it is obtained by a translation and a rotation from $x_1, x_2, ..., x_m$. We define a certain notion of convexity for Ω as follows.

Definition 4.1. We call $\Omega \subset R^m$ parabolically convex or p-convex in short if for any $p \in \partial\Omega$, there exists an orthonormal coordinate system $(\tilde{x}_1, \tilde{x}_2, ..., \tilde{x}_m)$ with the origin at p and $\Omega \subset \{a\tilde{x}_1^2 - \tilde{x}_2 < 0\}$ where $a > 0$ is independent of p.

Note that a C^2-smooth bounded domain with positively curved (positive principal curvatures) boundary is p-convex.

Proposition 4.4. *Let $\Omega \subset R^m$ be a p-convex bounded domain. Suppose $u_\varepsilon \in C^2(\Omega) \cap C^1(\bar{\Omega})$ satisfies $Q_\varepsilon u_\varepsilon = 0$ in Ω and $u_\varepsilon = \sigma\varphi \in C^2(\bar{\Omega})$ on $\partial\Omega$ with $\vec{F} \in C^1(\bar{\Omega})$ for $0 \le \sigma \le 1$. Then there exists a constant $C = C(\varepsilon, a, ||F_I||_\infty, ||\partial_I F_J||_\infty, ||\partial_I\varphi||_\infty, ||\partial_I\partial_J\varphi||_\infty)$ (independent of σ) such that*

$$\sup_{\partial\Omega} |\nabla u_\varepsilon| \le C. \tag{4.17}$$

Moreover, the bounds hold uniformly for $0 < \varepsilon \le \varepsilon_0$, a positive constant.

Proof. Given $p \in \partial\Omega$, we have an orthonormal coordinate system $(\tilde{x}_1, \tilde{x}_2, ..., \tilde{x}_m)$ as in the definition of p-convexity. Consider the comparison function $w = \alpha G + \sigma\varphi$ where G is the function $\tilde{G} \equiv a\tilde{x}_1^2 - \tilde{x}_2$ viewed as a function of (x_I), $I = 1, 2, ..., m$, for large α to be determined. In view of the invariance of $Q_\varepsilon(u)$ under the coordinate changes of translations and rotations, we compute ($\tilde{Q}_\varepsilon$, $\tilde{D}_\varepsilon$ being the corresponding operator, quantity of Q_ε, D_ε with respect to $(\tilde{x}_I)$, respectively)

$$\begin{aligned} Q_\varepsilon(w) &= \tilde{Q}_\varepsilon(\tilde{w}) \ \ (\tilde{w} \text{ is } w \text{ viewed as a function of } (\tilde{x}_I)) \\ &= \frac{P(\tilde{G})\alpha^3 + A\alpha^2 + B\alpha + E}{\tilde{D}_\varepsilon(\tilde{w})^3} \end{aligned} \tag{4.18}$$

by (4.2) where $P(\tilde{G})$ is the corresponding quantity of $P(G) \equiv G_{x_1}^2 G_{x_2x_2} - 2G_{x_1}G_{x_2}\, G_{x_1x_2} + G_{x_2}^2 G_{x_1x_1}$ with respect to $(\tilde{x}_I)$, and A is a function of $a,\ F_I,\ \partial_I F_J,\ \partial_I\varphi,\ \partial_I\partial_J\varphi$ while B, E are functions of $\varepsilon,\ a,\ F_I,\ \partial_I F_J,\ \partial_I\varphi,\ \partial_I\partial_J\varphi$. Moreover, a direct computation shows that $P(\tilde{G}) = 2a$. Since $a > 0$, $Q_\varepsilon(w) \geq$ ($\leq$, respectively) $0 = Q_\varepsilon(u_\varepsilon)$ for positive (negative, respectively) large $\alpha = \alpha(\varepsilon,\ a,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty,\ ||\partial_I\varphi||_\infty,\ ||\partial_I\partial_J\varphi||_\infty)$ by (4.18). Note that α is independent of σ and independent of ε for $0 < \varepsilon \leq \varepsilon_0$. On the other hand, $w = \alpha G + \sigma\varphi \leq$ ($\geq$, respectively) $\sigma\varphi = u_\varepsilon$ on $\partial\Omega$ since $G \leq 0$ on $\bar{\Omega}$ by the p-convexity. Therefore $w \leq$ ($\geq$, respectively)u_ε in Ω by the comparison principle for second order quasilinear operators (see e.g. Theorem 10.1 in [9]). Noting that $G(p) = 0$ and hence $w(p) = \sigma\varphi(p) = u_\varepsilon(p)$, we then have

$$\frac{\partial u_\varepsilon}{\partial \nu} \leq (\geq,\ \text{respectively}) \frac{\partial w}{\partial \nu} \tag{4.19}$$

where $\nu = -\partial_{\tilde{x}_2}$ at p. Observe that $\frac{\partial w}{\partial \nu}$ (for either positive or negative α) is bounded by a constant depending on $\varepsilon,\ a,\ ||F_I||_\infty,\ ||\partial_I F_J||_\infty,\ ||\partial_I\varphi||_\infty,\ ||\partial_I\partial_J\varphi||_\infty$, but independent of σ, p (moreover, the bounds hold for $0 < \varepsilon \leq \varepsilon_0$, a positive constant), so is $\frac{\partial u_\varepsilon}{\partial \nu}$ by (4.19). Since $u_\varepsilon = \sigma\varphi$ on $\partial\Omega$ and $u_\varepsilon - \sigma\varphi \in C^1(\bar{\Omega})$, we can easily show that the derivatives of $u_\varepsilon - \sigma\varphi$ in the $\tilde{x}_1$, $\tilde{x}_3$, ..., $\tilde{x}_m$) (except $\tilde{x}_2$) directions all vanish at p. It follows that in the $\tilde{x}_j$ ($j \neq 2$) direction, the derivative of u_ε is the same as the derivative of $\sigma\varphi$. So of course it is bounded by $||\nabla\varphi||_\infty$ (note that $0 \leq \sigma \leq 1$). Altogether we have proved (4.17).

Q.E.D.

Proof of Theorem A.

In order to apply Theorem 11.8 in [9] to solve the Dirichlet problem (4.1), we consider a family of equations:

$$\begin{aligned} Q_{\varepsilon,\sigma} u &\equiv div \frac{\nabla u + \sigma\vec{F}}{\sqrt{\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2}} = 0 \text{ in } \Omega \\ u &= \sigma\varphi \ \text{ on } \partial\Omega, \ \ 0 \leq \sigma \leq 1. \end{aligned} \tag{4.20}$$

Express $Q_{\varepsilon,\sigma} u = a_{IJ}(\varepsilon, x, \nabla u; \sigma)u_{IJ} + b(\varepsilon, x, \nabla u; \sigma)$ where $a_{IJ}(\varepsilon, x, \nabla u; \sigma)$ and $b(\varepsilon, x, \nabla u; \sigma)$ are given by (4.3) with $\vec{F}$ replaced by $\sigma\vec{F}$. It is then easy to check that the conditions (i), (ii), (iii) on page 287 of [9] are satisfied. To have an apriori Hölder estimate for ∇u, we invoke Theorem 13.2 in [9]. Comparing (4.20) with (13.2) in [9] gives

$$\mathbf{A}(x, u, \nabla u) = \frac{\nabla u + \sigma\vec{F}}{\sqrt{\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2}}, B(x, u, \nabla u) = 0.$$

Following pages 319-320 of [9], we find $\bar{a}^{IJ} \equiv D_{p_J}A^I = a_{IJ}(\varepsilon, x, \nabla u; \sigma)$ and $\lambda(\varepsilon, x, u, \nabla u) = \varepsilon^2/[\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2]^{3/2}$ by (4.4). Therefore we can take $\lambda_K = \varepsilon^2/[\varepsilon^2 + (K + C)^2]^{3/2}$ in (13.4) of [9], in which $K \equiv |u|_{1;\Omega}$ (see page 53 in [9] for the notation) and $C \equiv ||\vec{F}||_\infty$. Similarly we estimate

$$|D_{p_J}A^I| = |a_{IJ}(\varepsilon, x, \nabla u; \sigma)| \leq \frac{1}{\sqrt{\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2}} \leq \frac{1}{\varepsilon}.$$

So we can take $\Lambda_K = \varepsilon^{-1}$. Since both D_zA^I and B vanish, we compute

$$\begin{aligned} |\delta_J A^I| + |B| &= |D_{x_J}A^I| \\ &= \frac{|(\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2)\partial_J(\sigma F_I) - (u_I + \sigma F_I)(u_L + \sigma F_L)\partial_J(\sigma F_L)|}{[\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2]^{3/2}} \\ &\leq \frac{(\frac{3}{2} + n)\sup_{K,I}|\partial_K F_I|}{\sqrt{\varepsilon^2 + |\nabla u + \sigma\vec{F}|^2}}. \end{aligned}$$

Therefore we can take an upper bound $\mu_K = \varepsilon^{-1}(\frac{3}{2} + n)\sup_{J,I}|\partial_J F_I|$. Now by Theorem 13.2 in [9], we have an apriori Hölder bound for ∇u in terms of n, K $(\equiv |u|_{1;\Omega})$, Λ_K/λ_K, μ_K/λ_K, size of Ω, and $|\varphi|_{2;\Omega}$. On the other hand, we observe that Lemma 4.1, Propositions 4.2-4.4 still hold for $Q_{\varepsilon,\sigma}$ instead of Q_ε. So we have an apriori C^1 bound for solutions of (4.20), independent of σ and ε (for $0 < \varepsilon \leq \varepsilon_0$). Altogether we have obtained an apriori $C^{1,\beta}(\bar{\Omega})$ $(\beta > 0)$ bound for solutions of (4.20), independent of σ (but depend on ε). By Theorem 11.8 in [9], we obtain

Theorem 4.5. *Let Ω be a p-convex bounded domain in $R^m, m \geq 2$, with $\partial\Omega \in C^{2,\alpha}$ $(0 < \alpha < 1)$. Let $\varphi \in C^{2,\alpha}(\bar{\Omega})$. Suppose $\vec{F} \in C^{1,\alpha}(\bar{\Omega})$ satisfies the condition (4.11) for $C^{1,\alpha}$-smooth and bounded f_K's in Ω. Then there exists a solution $u_\varepsilon \in C^{2,\alpha}(\bar{\Omega})$ of the Dirichlet problem: $Q_\varepsilon(u) = 0$ in Ω, $u = \varphi$ on $\partial\Omega$ for given $\varepsilon > 0$.*

(Proof of Theorem A Continued)

Propositions 4.2-4.4 tell us that there exists a constant $C = C(\varepsilon,\ a,\ R,\ ||F_I||_\infty, ||\partial_I F_J||_\infty, ||\varphi||_\infty, ||\partial_I\varphi||_\infty, ||\partial_I\partial_J\varphi||_\infty, ||f_I||_\infty)$ such that

$$\sup_\Omega |u_\varepsilon| + \sup_\Omega |\nabla u_\varepsilon| \leq C. \tag{4.21}$$

Moreover, the bounds hold uniformly for $0 < \varepsilon \leq \varepsilon_0$, a positive constant.

In view of (4.21) we can find a subsequence u_{ε_j} $(0 < \varepsilon_j \leq \varepsilon_0,\ \varepsilon_j \to 0)$ converging to u_0 in C^0 by the Arzela-Ascoli theorem. Then the Lipschitzianity of u_0 follows by taking the limit of ratios: $(x \neq y)$

$$\left|\frac{u_{\varepsilon_j}(x) - u_{\varepsilon_j}(y)}{x - y}\right| \ (\leq C).$$

Next we claim that u_0 is a minimizer for $\mathcal{F}(\cdot)$ (see (1.3)) such that $u_0 = \varphi$ on $\partial\Omega$. Observe that $W^{1,q}(\Omega)$ is compactly imbedded in $L^1(\Omega)$ (e.g., Theorem 7.26 in [9]). So we may as well assume that u_{ε_j} converges to u_0 in $L^1(\Omega)$. Also note that $|\vec{p} + \vec{F}|$ is convex in $\vec{p}$ since $|\lambda\vec{p}_1 + (1-\lambda)\vec{p}_2 + \vec{F}| = |\lambda(\vec{p}_1 + \vec{F}) + (1-\lambda)(\vec{p}_2 + \vec{F})|$

$\leq \lambda|\vec{p}_1+\vec{F}| + (1-\lambda)|\vec{p}_2+\vec{F}|$ for $0 \leq \lambda \leq 1$. We can therefore apply Theorem 4.1.2 in [15] to conclude the lower semicontinuity of $\mathcal{F}(\cdot)$ (see (1.3)):

$$\mathcal{F}(u_0) \leq \liminf_{j\to\infty} \mathcal{F}(u_{\varepsilon_j}). \tag{4.22}$$

Now for $v \in W^{1,1}$ with $v - \varphi \in W_0^{1,1}$, we estimate

$$\begin{aligned} \mathcal{F}(u_{\varepsilon_j}) &\equiv \int_\Omega |\ \nabla u_{\varepsilon_j} + \vec{F}\ | \ \text{ (omitting volume element)} \\ &\leq \int_\Omega \sqrt{\varepsilon_j^2 + |\nabla u_{\varepsilon_j} + \vec{F}|^2} \\ &\leq \int_\Omega \sqrt{\varepsilon_j^2 + |\nabla v + \vec{F}|^2} \\ &\leq \varepsilon_j\ vol(\Omega) + \int_\Omega |\nabla v + \vec{F}| \end{aligned} \tag{4.23}$$

where we have used the fact that the Dirichlet solution $u_{\varepsilon_j} \in C^2(\bar{\Omega})$ is also a minimizer for $\mathcal{F}_{\varepsilon_j}(u) \equiv \int_\Omega \sqrt{\varepsilon_j^2 + |\nabla u + \vec{F}|^2}$. Taking the limit infimum of (4.23) and making use of (4.22), we finally obtain that $\mathcal{F}(u_0) \leq \int_\Omega |\nabla v + \vec{F}| \equiv \mathcal{F}(v)$. That is to say, u_0 is a minimizer for $\mathcal{F}(\ \cdot\ )$.

Q.E.D.

5. Uniqueness of minimizers-proof of Theorems B and C

Recall (see Section 3) that $\Omega \subset R^m$ denotes a bounded domain and $\mathcal{F}(u) \equiv \int_\Omega \{|\nabla u + \vec{F}| + Hu\}$ for $u \in W^{1,1}(\Omega)$, $\vec{F} \in L^1(\Omega)$, and $H \in L^\infty(\Omega)$. We will prove two ($W^{1,1}$) minimizers for $\mathcal{F}(u)$ with the same "boundary value" have the same normal vector "almostly".

Theorem 5.1. *Let $u, v \in W^{1,1}(\Omega)$ be two minimizers for $\mathcal{F}(u)$ such that $u - v \in W_0^{1,1}(\Omega)$. Let $u_\varepsilon \equiv u + \varepsilon(v-u)$. Then for any pair of regular ε_1, $\varepsilon_2 \in [0,1]$, there holds $N(u_{\varepsilon_1}) = N(u_{\varepsilon_2})$ in $\Omega \backslash [S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$(a.e.).*

Proof. By (3.13) with $\varphi = v - u$, we have

$$0 = \mathcal{F}(v) - \mathcal{F}(u) = \int_0^1 \frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} d\varepsilon. \tag{5.1}$$

As in the proof of Theorem 3.3, the same argument shows that $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} \geq 0$ for any regular $\varepsilon \in [0,1]$. In view of (5.1) and Lemma 3.2(1), $\frac{d\mathcal{F}(u_\varepsilon)}{d\varepsilon} = 0$ for any regular $\varepsilon \in [0,1]$. It follows from (3.4) that $\int_{\Omega\backslash S(u_\varepsilon)} N(u_\varepsilon)\cdot\nabla(v-u) = 0$. Therefore for any pair of regular ε_1, $\varepsilon_2 \in [0,1]$, there holds

$$\int_{\Omega\backslash[S(u_{\varepsilon_1})\cup S(u_{\varepsilon_2})]} [N(u_{\varepsilon_2}) - N(u_{\varepsilon_1})]\cdot\nabla(v-u) = 0. \tag{5.2}$$

Here we have used $\int_{S(u_{\varepsilon_1})\backslash S(u_{\varepsilon_2})} N(u_{\varepsilon_2}) \cdot \nabla(v-u) = 0$ and $\int_{S(u_{\varepsilon_2})\backslash S(u_{\varepsilon_1})} N(u_{\varepsilon_1}) \cdot \nabla(v-u) = 0$ by observing that for $j = 1, 2$, $|N(u_{\varepsilon_j}) \cdot \nabla(v-u)| \leq |\nabla(v-u)|$ and $\int_{S(u_{\varepsilon_j})} |\nabla(v-u)| = 0$ from the definition of ε_j being regular. Write $v - u = (u_{\varepsilon_2} - u_{\varepsilon_1})/(\varepsilon_2 - \varepsilon_1)$ for $\varepsilon_2 \neq \varepsilon_1$. By Lemma 5.1' in [3], the integrand in (5.2) is

$$\frac{|\nabla u_{\varepsilon_2} + \vec{F}| + |\nabla u_{\varepsilon_1} + \vec{F}|}{2(\varepsilon_2 - \varepsilon_1)} |N(u_{\varepsilon_2}) - N(u_{\varepsilon_1})|^2.$$

It then follows that $N(u_{\varepsilon_1}) = N(u_{\varepsilon_2})$ in $\Omega\backslash[S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$.

Q.E.D.

For a vector field $\vec{G} = (g_1, g_2, ..., g_{2n})$ on $\Omega \subset R^{2n}$, we recall that $\vec{G}^* \equiv (g_2, -g_1, g_4, -g_3, ..., g_{2n}, -g_{2n-1})$.

Lemma 5.2. *Let $u, v \in W^1(\Omega)$ where the domain Ω is contained in R^{2n}. Let $u_\varepsilon \equiv u + \varepsilon(v-u)$. Suppose $N(u_{\varepsilon_1}) = N(u_{\varepsilon_2})$ in $\Omega\backslash[S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$ for a pair ε_1, ε_2 such that $\varepsilon_1 \neq \varepsilon_2$. Then for $j = 1, 2$, there holds*

$$(\nabla u_{\varepsilon_j} + \vec{F})^* \cdot (\nabla v - \nabla u) = 0 \text{ in } \Omega \text{ (a.e.)}. \tag{5.3}$$

Proof. We will prove (5.3) only for $j = 1$ (similar argument works also for $j = 2$) For $p \in S(u_{\varepsilon_1})$, $\nabla u_{\varepsilon_1} + \vec{F} = 0$. So (5.3) holds obviously. For $p \in S(u_{\varepsilon_2})$, (5.3) also holds by observing that $\nabla v - \nabla u = [(\nabla u_{\varepsilon_1} + \vec{F}) - (\nabla u_{\varepsilon_2} + \vec{F})]/(\varepsilon_1 - \varepsilon_2) = (\nabla u_{\varepsilon_1} + \vec{F})/(\varepsilon_1 - \varepsilon_2)$ and $\vec{G}^* \cdot \vec{G} = 0$. For the remaining case: $p \in \Omega\backslash[S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$, we observe that for $j = 1, 2$,

$$N(u_{\varepsilon_j})^* \cdot \nabla u_{\varepsilon_j} = \frac{\vec{F}^* \cdot \nabla u_{\varepsilon_j}}{|\nabla u_{\varepsilon_j} + \vec{F}|} = \vec{F}^* \cdot N(u_{\varepsilon_j}). \tag{5.4}$$

Here we have used the property $\vec{G}^* \cdot \vec{G} = 0$ twice. Since $N(u_{\varepsilon_1}) = N(u_{\varepsilon_2})$ in $\Omega\backslash[S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$ by assumption (hence $N(u_{\varepsilon_1})^* = N(u_{\varepsilon_2})^*$ also), we take the difference of (5.4) for $j = 1$ and $j = 2$ to obtain

$$N(u_{\varepsilon_1})^* \cdot (\nabla u_{\varepsilon_2} - \nabla u_{\varepsilon_1}) = 0. \tag{5.5}$$

Formula (5.3) for $j = 1$ on $\Omega\backslash[S(u_{\varepsilon_1}) \cup S(u_{\varepsilon_2})]$ then follows from (5.5) by noting that $v - u = (u_{\varepsilon_2} - u_{\varepsilon_1})/(\varepsilon_2 - \varepsilon_1)$.

Q.E.D.

We will use the following general criterion to prove the uniqueness of minimizers and a comparison principle for weak functions later.

Theorem 5.3. *Let Ω be a bounded domain in R^{2n}. Let $w \in W_0^{1,p}(\Omega)$, $\sigma \in W^{1,q}(\Omega)$, where $1 \leq p < \infty$, $q = \frac{p}{p-1}$ ($q = \infty$ for $p = 1$). Let $\vec{F}$ (a vector field) $\in W^{1,1}(\Omega) \cap L^q(\Omega)$ satisfying $div\vec{F}^* > 0$ (a.e.) or $div\vec{F}^* < 0$ (a.e.). Suppose $(\nabla\sigma + \vec{F})^* \cdot \nabla w = 0$ in Ω (a.e.). Then $w \equiv 0$ in Ω (a.e.).*

Proof. Take $\omega_j \in C_0^\infty(\Omega) \to w$ in $W^{1,p}$ and $\vec{F}_{\bar{k}} \in C^\infty(\Omega) \to \vec{F}$ in $W^{1,1} \cap L^q$. Suppose ω_j does not vanish identically. Then there exists a decreasing sequence of positive numbers a_i converging to 0 such that $\Omega_{j,i} \equiv \{|\omega_j| > a_i\} \subset\subset \Omega$ is not empty for large i and $\partial\Omega_{j,i}$ is C^∞ smooth (by Sard's theorem; note that $|\omega_j|$ is C^∞ smooth where $\omega_j \neq 0$). Also we take $v_k \in C^\infty(\Omega) \to \sigma$ in $W^{1,2}$. Consider

$$I_{j,i,k,\bar{k}} \equiv \int_{\partial\Omega_{j,i}} |\omega_j| \; (\nabla v_k + \vec{F}_{\bar{k}})^* \cdot \nu \tag{5.6}$$

where ν denotes the boundary normal. We first compute

$$\begin{aligned} \int_{\partial\Omega_{j,i}} |\omega_j| \; (\nabla v_k + \vec{F}_{\bar{k}})^* \cdot \nu &= a_i \int_{\partial\Omega_{j,i}} (\nabla v_k + \vec{F}_{\bar{k}})^* \cdot \nu \\ &= a_i \int_{\Omega_{j,i}} div[(\nabla v_k)^* + \vec{F}_{\bar{k}}^*] \\ &= a_i \int_{\Omega_{j,i}} div \vec{F}_{\bar{k}}^* \end{aligned} \tag{5.7}$$

Here we have used Green's theorem for the second equality and $div(\nabla v_k)^* = 0$ for the third equality in (5.7). It follows from (5.7) that

$$\lim_{i \to \infty} I_{j,i,k,\bar{k}} = 0. \tag{5.8}$$

On the other hand, a similar reasoning gives

$$\begin{aligned} I_{j,i,k,\bar{k}} &= \int_{\Omega_{j,i}} \nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div[(\nabla v_k)^* + \vec{F}_{\bar{k}}^*] \\ &= \int_{\Omega_{j,i}} \nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div\vec{F}_{\bar{k}}^*. \end{aligned} \tag{5.9}$$

Observe that $\cup_i \Omega_{j,i} = \{|\omega_j| > 0\} = \Omega \backslash \{\omega_j = 0\}$, $(\Omega\backslash\{\omega_j = 0\})\backslash\Omega_{j,i} = \cup_{l=i}^\infty (\Omega_{j,l+1} \backslash \Omega_{j,l})$, $\vec{F}_{\bar{k}} \in W^{1,1}(\Omega)$, and hence

$$\begin{aligned} &(\int_{\Omega_{j,i}} - \int_{\Omega\backslash\{\omega_j=0\}})\{\nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div\vec{F}_{\bar{k}}^*\} \\ = \;& -\Sigma_{l=i}^\infty \int_{\Omega_{j,l+1}\backslash\Omega_{j,l}} \{\nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div\vec{F}_{\bar{k}}^*\} \\ = \;& -\Sigma_{l=i}^\infty \; (I_{j,l+1,k,\bar{k}} - I_{j,l,k,\bar{k}}) = I_{j,i,k,\bar{k}} \end{aligned} \tag{5.10}$$

by (5.8). It follows from (5.9), (5.10) that

$$\begin{aligned} 0 &= \int_{\Omega\backslash\{\omega_j=0\}} \nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div\vec{F}_{\bar{k}}^* \\ &= \int_\Omega \nabla|\omega_j| \cdot (\nabla v_k + \vec{F}_{\bar{k}})^* + |\omega_j| \; div\vec{F}_{\bar{k}}^*. \end{aligned}$$

Here we have used $\nabla|\omega_j| = 0$ if $\omega_j = 0$ (p.152 in [9]). Letting $\bar{k} \to \infty$ in the above formula gives

$$0 = \int_\Omega \nabla|\omega_j| \cdot (\nabla v_k + \vec{F})^* + |\omega_j| \; div\vec{F}^*. \tag{5.11}$$

Letting $k \to \infty$ in the first term of (5.11), we then estimate by using the assumption $\nabla w \cdot (\nabla\sigma + \vec{F})^* = 0$

$$\begin{aligned} &\int_\Omega \nabla|\omega_j| \cdot (\nabla\sigma + \vec{F})^* \\ = &\int_{\{\omega_j>0\}} (\nabla\omega_j - \nabla w) \cdot (\nabla\sigma + \vec{F})^* - \int_{\{\omega_j<0\}} (\nabla\omega_j - \nabla w) \cdot (\nabla\sigma + \vec{F})^* \\ \longrightarrow \;& 0 \quad as \;\; j \to \infty. \end{aligned} \tag{5.12}$$

Here we have used $\omega_j \to w$ in $W^{1,p}$ and $(\nabla\sigma + \vec{F})^* \in L^q(\Omega)$ by assumption. For the second term of (5.11), we have

$$\lim_{j\to\infty} \int_\Omega |\omega_j| \; div\vec{F}^* = \int_\Omega |w| \; div\vec{F}^* > 0 \text{ or } < 0 \tag{5.13}$$

if $w \neq 0$ (noting that $div\vec{F}^* > 0$ or < 0 by assumption). By (5.11), (5.12), and (5.13), we reach a contradiction. Therefore $w \equiv 0$ in Ω (a.e.).

Q.E.D.

Remark. If $\vec{F}$ does not satisfy the condition in Theorem 5.3, then the theorem may not hold as shown by the following examples. Let $\Omega = (0,\pi) \times (0,\pi) \subset R^2$. Let $w = \sin x \sin y \in W_0^{1,2}$. Then $\nabla w = (\cos x \sin y, \, \sin x \cos y)$. Take $\sigma = 0$ and $\vec{F} = (\cos x \sin y, \, \sin x \cos y)$. It is easy to see that $\vec{F}^* = (\sin x \cos y, \, -\cos x \sin y)$, $div\vec{F}^* = 0$, and $(\nabla\sigma + \vec{F})^* \cdot \nabla w = \vec{F}^* \cdot \nabla w = 0$. With the same σ $(= 0)$ and w as above, we can also take $\vec{F} = \sin x \; (\cos x \sin y, \sin x \cos y)$. Then still $(\nabla\sigma + \vec{F})^* \cdot \nabla w$ $= \vec{F}^* \cdot \nabla w = 0$ while $div\vec{F}^* = \cos x \; \sin x \; \cos y$ has no definite sign in Ω.

Proof of Theorem B.

The proof follows from Theorem 5.1, Lemma 5.2, and Theorem 5.3 with $p = q = 2$, $\sigma = u_{\varepsilon_1}$, and $w = v - u$.

Q.E.D.

Next we want to prove a comparison principle for weak sub- and super- solutions (a comparison principle for C^2-smooth functions has been studied in [3]. See Theorem C and Theorem C' there). First we need to define relevant differential inequalities in some weak sense. Let $\Omega \subset R^m$ denote a bounded domain. Recall that $N(u) \equiv \frac{\nabla u + \vec{F}}{|\nabla u + \vec{F}|}$ is defined on $\Omega \backslash S(u)$ ($\vec{F}$, say, is an L^1_{loc} vector field in Ω).

Definition 5.1. Let $H \in L^1_{loc}(\Omega)$. We say $u \in W^{1,1}(\Omega)$ satisfies $divN(u) \geq H$ ($\leq H$, respectively) in the weak sense in Ω if and only if for any $\varphi \in C_0^\infty(\Omega)$ and $\varphi \geq 0$, there holds

$$-\int_{S(u)}|\nabla\varphi|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi+\int_{\Omega}H\varphi \quad\leq\quad 0 \tag{5.14}$$

$$(\int_{S(u)}|\nabla\varphi|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi+\int_{\Omega}H\varphi \quad\geq\quad 0,\ \text{respectively}). \tag{5.15}$$

Recall that we defined the weak solution to $divN(u) = H$ in Section 3 (see (3.12)). The following result justifies the above definitions.

Proposition 5.4. *Let $H \in L^{\infty}(\Omega)$. Then $u \in W^{1,1}(\Omega)$ satisfies $divN(u) \geq H$ and $divN(u) \leq H$ in the weak sense if and only if $u \in W^{1,1}(\Omega)$ is a weak solution to the equation $divN(u) = H$.*

Proof. Since $C_0^{\infty}(\Omega)$ is dense in $W_0^{1,1}(\Omega)$, (5.14) and (5.15) hold for every $\varphi \in C_0^{\infty}(\Omega)$ if and only if they hold for every $\varphi \in W_0^{1,1}(\Omega)$. Write $\varphi = \varphi^{+} - \varphi^{-}$ for $\varphi \in W_0^{1,1}(\Omega)$ where $\varphi^{+} \equiv \max\{\varphi, 0\}$ and $\varphi^{-} \equiv \max\{-\varphi, 0\}$. Express

$$\begin{aligned} & \int_{S(u)}|\nabla\varphi|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi+\int_{\Omega}H\varphi \\ = \ & \{\int_{S(u)}|\nabla\varphi^{+}|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi^{+}+\int_{\Omega}H\varphi^{+}\} \\ & -\{-\int_{S(u)}|\nabla\varphi^{-}|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi^{-}+\int_{\Omega}H\varphi^{-}\}. \end{aligned} \tag{5.16}$$

Note that $\varphi^{+} \geq 0$ and $\varphi^{-} \geq 0$. Now suppose u is a weak solution to $divN(u) \leq H$ *and* $divN(u) \geq H$. Then the right-hand side of (5.16) is nonnegative by our definitions. So the left hand side of (5.16) is nonnegative, i.e., (3.12) holds. Conversely, suppose u is a weak solution to $divN(u) = H$. That is to say, the left hand side of (5.16) is nonnegative (note that φ is not restricted to be nonnegative here). By taking $\varphi \geq 0$ i.e. $\varphi^{-} = 0$ ($\varphi \leq 0$ i.e. $\varphi^{+} = 0$,respectively) in (5.16), we obtain (5.15) ((5.14), respectively).

Q.E.D.

Definition 5.2. $u, v \in W^{1}(\Omega)$ satisfy $divN(u) \geq divN(v)$ in Ω in the weak sense if and only if for any $\varphi \in W_0^{1,1}(\Omega)$ and $\varphi \geq 0$, there holds

$$-\int_{S(u)}|\nabla\varphi|+\int_{\Omega\backslash S(u)}N(u)\cdot\nabla\varphi \leq +\int_{S(v)}|\nabla\varphi|+\int_{\Omega\backslash S(v)}N(v)\cdot\nabla\varphi. \tag{5.17}$$

Definition 5.3. $u, v \in W^{1,1}(\Omega)$ satisfy $u \leq v$ on $\partial\Omega$ if and only if $(u-v)^{+} \equiv \max(u-v, 0) \in W_0^{1,1}(\Omega)$.

Theorem 5.5. *Suppose $u, v \in W^{1,1}(\Omega)$ satisfy the following conditions:*

$$\begin{aligned} divN(u) &\geq divN(v) \text{ in } \Omega \text{ (in the weak sense);} \\ u &\leq v \text{ on } \partial\Omega. \end{aligned}$$

Then $N(u) = N(v)$ on $\{u > v\}\backslash[S(u) \cup S(v)]$.

Proof. Let $\varphi = (u-v)^+$. The condition $u \leq v$ on $\partial\Omega$ implies that $\varphi \in W_0^{1,1}(\Omega)$. Let $v_\varepsilon \equiv v + \varepsilon\varphi$. From Lemma 3.2 (1), $\frac{d\mathcal{F}(v_\varepsilon)}{d\varepsilon}$ is increasing in regular ε. It follows that $\frac{d\mathcal{F}(v_{0+})}{d\varepsilon} \leq \frac{d\mathcal{F}(v_{1-})}{d\varepsilon}$ by Lemma 3.2 (2). In view of the formula (3.3), we have

$$+\int_{S(v)} |\nabla\varphi| + \int_{\Omega\backslash S(v)} N(v)\cdot\nabla\varphi \leq -\int_{S(v_1)} |\nabla\varphi| + \int_{\Omega\backslash S(v_1)} N(v_1)\cdot\nabla\varphi. \tag{5.18}$$

Observe that $v_1 = u$ on $\{u > v\}$ and $\varphi = 0$ on $\{u \leq v\}$. So the right hand side of (5.18) equals the left hand side of (5.17). It follows that

$$-\int_{S(u)} |\nabla\varphi| + \int_{\Omega\backslash S(u)} N(u)\cdot\nabla\varphi = +\int_{S(v)} |\nabla\varphi| + \int_{\Omega\backslash S(v)} N(v)\cdot\nabla\varphi. \tag{5.19}$$

Write

$$\begin{aligned}
&\int_{\Omega\backslash S(u)} N(u)\cdot\nabla\varphi - \int_{\Omega\backslash S(v)} N(v)\cdot\nabla\varphi \\
=\ &\int_{\Omega\backslash[S(u)\cup S(v)]} (N(u)-N(v))\cdot\nabla\varphi \\
&+\int_{S(v)\backslash S(u)} N(u)\cdot\nabla\varphi - \int_{S(u)\backslash S(v)} N(v)\cdot\nabla\varphi.
\end{aligned} \tag{5.20}$$

We claim

$$-\int_{S(u)} |\nabla\varphi| - \int_{S(u)\backslash S(v)} N(v)\cdot\nabla\varphi = 0. \tag{5.21}$$

Since $\varphi = 0$ on $\{u \leq v\}$, we only have to discuss the case that $u > v$. In this case, $\varphi = u - v$ and hence $\nabla\varphi = (\nabla u + \vec{F}) - (\nabla v + \vec{F}) = -(\nabla v + \vec{F})$ in $S(u)$ (and $= 0$ in $S(u)\cap S(v)$). So $N(v)\cdot\nabla\varphi = \frac{\nabla v+\vec{F}}{|\nabla v+\vec{F}|}\cdot[-(\nabla v + \vec{F})] = -|\nabla v + \vec{F}|$ in $S(u)\backslash S(v)$. It is now clear that (5.21) holds. Similarly there also holds

$$-\int_{S(v)} |\nabla\varphi| + \int_{S(v)\backslash S(u)} N(u)\cdot\nabla\varphi = 0. \tag{5.22}$$

Combining (5.19), (5.20), (5.21), (5.22) gives

$$\int_{\Omega\backslash[S(u)\cup S(v)]} (N(u)-N(v))\cdot\nabla\varphi = 0. \tag{5.23}$$

By Lemma 5.1' in [3] (which works also for $u, v \in W^{1,1}(\Omega)$), we have

$$(N(u)-N(v))\cdot\nabla\varphi = \frac{|\nabla u+\vec{F}| + |\nabla v+\vec{F}|}{2}|N(u)-N(v)|^2 \tag{5.24}$$

on $\{u > v\}\backslash[S(u)\cup S(v)]$ (where $\varphi = u - v$). Noting that $\varphi = 0$ on $\{u \leq v\}$ and substituting (5.24) into (5.23), we finally obtain $N(u) = N(v)$ on $\{u > v\}\backslash[S(u)\cup S(v)]$.

Q.E.D.

We can now prove the comparison principle for weak sub- and super- solutions.

Proof of Theorem C.

By Theorem 5.5 and Lemma 5.2 (switching the roles of u and v and taking $\Omega = \{u > v\}$, $\varepsilon_1 = 0$, $\varepsilon_2 = 1$), we obtain $(\nabla v + \vec{F})^* \cdot \nabla(u-v)^+ = 0$. Then we apply Theorem 5.3 (with $p = q = 2$, $\sigma = v$, and $w = (u-v)^+$) to conclude that $(u-v)^+ = 0$ in Ω. That is to say, $u \leq v$ in Ω.

Q.E.D.

6. When a smooth solution is a minimizer

In this section we determine when a smooth solution is a minimizer. We will prove Theorem D, Theorem E, and Corollary F. We first prove a result for the case $H_{m-1}(S(u)) = 0$, in which a C^2-smooth solution must be a weak solution.

Lemma 6.1. *Let Ω be a bounded domain in R^m. Suppose $u \in C^1(\Omega) \cap C^2(\Omega \backslash S(u)) \cap C^0(\bar{\Omega})$ satisfies (1.6) in $\Omega \backslash S(u)$ with $\vec{F} \in C^1(\Omega \backslash S(u))$ and $H \in C^0(\Omega \backslash S(u)) \cap L^1_{loc}(\Omega)$. Suppose $H_{m-1}(S(u))$, the $m-1$ dimensional Hausdorff measure of $S(u)$, vanishes. Then u is a weak solution to (1.6) and a minimizer for (1.7) if $u \in W^{1,1}(\Omega)$ and $H \in L^\infty(\Omega)$ also.*

Proof. By Theorem 3.3, it suffices to prove that for any $\varphi \in C_0^\infty(\Omega)$ (3.12) holds. That is,

$$\int_{S(u)} |\nabla\varphi| + \int_{\Omega\backslash S(u)} N(u) \cdot \nabla\varphi + \int_\Omega H\varphi \geq 0.$$

Write $\Omega = \Omega_+ \cup \Omega_0 \cup \Omega_-$ where $\Omega_+ \equiv \{\varphi > 0\}$, $\Omega_- \equiv \{\varphi < 0\}$, and $\Omega_0 \equiv \{\varphi = 0\}$. If $\Omega_+ \neq \emptyset$, then there exists a sequence of $\varepsilon_j > 0$ approaching 0, such that $\Omega_{\varepsilon_j} \equiv \{\varphi > \varepsilon_j\} \neq \emptyset$, $\cup_{j=1}^\infty \Omega_{\varepsilon_j} = \Omega_+$ and $\partial\Omega_{\varepsilon_j}$ are C^∞-smooth by Sard's theorem. Since $u \in C^1(\Omega)$, $S(u) \cap \bar{\Omega}_{\varepsilon_j}$ is compact. Together with the condition $H_{m-1}(S(u)) = 0$, for any $\alpha > 0$, we can find a finite cover of balls $B_{r_k}(p_k)$ of center p_k and radius r_k, $k = 1, 2, \ldots, K$ for $S(u) \cap \bar{\Omega}_{\varepsilon_j}$ such that

$$\sum_{k=1}^K H_{m-1}(\partial B_{r_k}(p_k)) < \alpha. \tag{6.1}$$

On the other hand we compute by the divergence theorem and the equation (1.6)

$$\begin{aligned} &\int_{\partial(\Omega_{\varepsilon_j}\backslash\cup B_{r_k}(p_k))} (\varphi - \varepsilon_j) N(u) \cdot \nu \\ = &\int_{\Omega_{\varepsilon_j}\backslash\cup B_{r_k}(p_k)} \nabla\varphi \cdot N(u) + (\varphi - \varepsilon_j) H. \end{aligned} \tag{6.2}$$

Since $\varphi - \varepsilon_j = 0$ on $\partial\Omega_{\varepsilon_j}$, we can estimate the boundary term in (6.2) as follows:

$$\begin{aligned}(6.3)\qquad & \Big|\int_{\partial(\Omega_{\varepsilon_j}\backslash\cup B_{r_k}(p_k))}(\varphi-\varepsilon_j)N(u)\cdot\nu\,\Big| \\ \leq\ & \{\max_{\Omega}|\varphi-\varepsilon_j|\}H_{m-1}(\cup_{k=1}^{K}\partial B_{r_k}(p_k)) \\ \leq\ & \alpha\max_{\Omega}|\varphi-\varepsilon_j|\end{aligned}$$

by (6.1) and the fact that $|N(u)| = |\nu| = 1$. Letting $\alpha \to 0$ in (6.3) gives

$$\int_{\Omega_{\varepsilon_j}\backslash S(u)}\nabla\varphi\cdot N(u)+(\varphi-\varepsilon_j)H=0 \tag{6.4}$$

in view of (6.2). Letting $\varepsilon_j \to 0$ in (6.4), we obtain

$$\int_{\Omega_+\backslash S(u)}\nabla\varphi\cdot N(u)+\int_{\Omega_+}\varphi H=0 \tag{6.5}$$

by noting that the volume of $\{0<\varphi\leq\varepsilon_j\}$ tends to 0 as $\varepsilon_j \to 0$. Similarly we also have

$$\int_{\Omega_-\backslash S(u)}\nabla\varphi\cdot N(u)+\int_{\Omega_-}\varphi H=0. \tag{6.6}$$

On the other hand, it is obvious that the integral of φH over Ω_0 vanishes since $\varphi = 0$ on Ω_0. Observing that $\nabla\varphi = 0$ a.e. on Ω_0 in view of Lemma 7.7 in [9], we conclude that

$$\int_{\Omega_0\backslash S(u)}\nabla\varphi\cdot N(u)=0. \tag{6.7}$$

It now follows from (6.5), (6.6), and (6.7) that

$$\int_{\Omega\backslash S(u)}\nabla\varphi\cdot N(u)+\int_{\Omega}\varphi H=0 \tag{6.8}$$

for $\varphi \in C_0^\infty(\Omega)$. Comparing (6.8) with (3.12) and noting that the first integral of (3.12) is zero by $H_{m-1}(S(u)) = 0$, we have completed the proof.

Q.E.D.

Proof of Theorem D.

Write $\nabla u + \vec{F} = (u_I + F_I)_{I=1}^m$. Consider the map $G : p \in \Omega \to ((u_I + F_I)(p))_{I=1}^m$. Computing the differential dG of G at a singular point p (where $G(p) = 0$), we obtain $(\partial_J u_I + \partial_J F_I)$ in matrix form (note that $G \in C^1$). From elementary linear algebra we compute

$$\begin{aligned}(6.9)\qquad & rank\ (\partial_J u_I+\partial_J F_I)+rank\ (\partial_I u_J+\partial_I F_J) \\ \geq\ & rank\ \{(\partial_J u_I+\partial_J F_I)-(\partial_I u_J+\partial_I F_J)\} \\ =\ & rank\ (\partial_J F_I-\partial_I F_J).\end{aligned}$$

Observing that $rank\ (\partial_J u_I + \partial_J F_I) = rank\ (\partial_I u_J + \partial_I F_J)$ (the transpose has the same rank), we can deduce from (6.9) that $rank\ dG(p) \geq [\frac{rank\ (h_{JI}(p))+1}{2}]$ where $h_{JI} \equiv (\partial_J F_I - \partial_I F_J)$. It follows that

$$dim(Ker\ dG(p)) \leq m - [\frac{rank\ (h_{JI}(p)) + 1}{2}]. \tag{6.10}$$

Then by the implicit function theorem there exists an open neighborhood V of p in Ω such that $G^{-1}(0) \cap V = S(u) \cap V$ is a submanifold of V, having (Euclidean) dimension dim_E bounded by the right side of (6.10).

Q.E.D.

Proof of Theorem E.

It suffices to prove that $H_{m-1}(S(u)) = 0$ in view of Lemma 6.1. Combining (1.9) and (1.10), we bound $dim_E S(u)$ by $m-2$. It follows that $H_{m-1}(S(u)) = 0$.

Q.E.D.

Proof of Corollary F.

For $m = 2n$, $\vec{F} = -\vec{X}^*$, we compute $rank\ (h_{JI}) = 2n$. Therefore (1.10) is reduced to $n \geq 2$, hence $m \geq 4$.

Q.E.D.

We remark that the condition (1.10) does not hold in dimension $m = 2$. So $H_1(S(u))$ may not vanish. Therefore a C^2-smooth solution may not be a minimizer in this case (see Example 7.4). We will discuss the general situation that $H_{m-1}(S(u)) > 0$ below.

First we will give a criterion for, in particular, a C^2-smooth solution to be a minimizer. Let Ω be a domain in R^m. Let $\Gamma \subset \Omega$ be a $m-1$ dimensional, orientable, C^1-smooth submanifold. Let $B \subset\subset \Omega$ be an open neighborhood of a point in Γ with C^1-smooth boundary and $\bar{B}$ being compact. Suppose $\Gamma \cap B$ divides B into two disjoint parts (note that Γ may or may not contain some singular points). That is, $B \backslash \Gamma = B \backslash\ (\Gamma \cap B) = B^+ \cup B^-$ where B^+ and B^- are disjoint domains (proper open and connected) (see Figure 1(a) or Figure 1(b) below). Suppose u is C^2-smooth in $\Omega \backslash \Gamma$ and has no singular points in $\Omega \backslash \Gamma$. Let $\vec{F} \in C^1(\Omega)$ for simplicity. Suppose also $N^+(u)$ and $N^-(u)$ (restrictions of $N(u)$ to B^+ and B^-, respectively) are continuous up to $\Gamma \cap B$, i.e., $N^+(u) \in C^0(\bar{B^+})$, $N^-(u) \in C^0(\bar{B^-})$, so that $divN^{\pm}(u) = H$ in $B^{\pm}$, respectively. Let ν^+ and ν^- denote the outward unit normals to $\Gamma \cap B$ with respect to B^+ and B^-, respectively. Note that $\nu^+ = -\nu^-$.

Proposition 6.2. *Suppose we have the situation described above. Then* u *is a weak solution to (1.6) on* B *with* $H \in C^0(B \backslash \Gamma) \cap L^\infty(B)$ *if and only if along* $\Gamma \cap B$, *there holds*

$$(N^+(u) - N^-(u)) \cdot \nu^+ = (N^+(u) - N^-(u)) \cdot \nu^- = 0. \tag{6.11}$$

Note that for $u \in W^{1,1}(B)$, u is a weak solution to (1.6) if and only if u is a minimizer for (1.7) in view of Theorem 3.3.

Proof. Using the divergence theorem, we compute



(a) (b)

FIGURE 1.

$$\begin{aligned}
(6.12) \qquad \int_{B\backslash\Gamma} N(u)\cdot\nabla\varphi + H\varphi &= (\int_{B^+} + \int_{B^-})(N(u)\cdot\nabla\varphi + H\varphi) \\
&= \int_{\partial B^+} \varphi N^+(u)\cdot\nu^+ + \int_{\partial B^-} \varphi N^-(u)\cdot\nu^- \\
&= \int_{\Gamma\cap B} \varphi(N^+(u) - N^-(u))\cdot\nu^+.
\end{aligned}$$

Here we have used $\nu^- = -\nu^+$ and $divN(u) = H$ in both B^+ and B^-. Observing that $H_m(S(u)\cap B) = 0$ since $H_m(S(u)\cap B) \leq H_m(\Gamma\cap B) = 0$, we conclude from (3.12) (also φ replaced by $-\varphi$) that u is a weak solution to (1.6) if and only if

$$(6.13) \qquad \int_{B\backslash\Gamma} N(u)\cdot\nabla\varphi + H\varphi = 0$$

for all $\varphi \in C_0^\infty(B)$. On the other hand, (6.13) holds if and only if (6.11) holds by (6.12).

Q.E.D.

In order to have a criterion for a more general situation, we extend Proposition 6.2 as follows. Let $\Omega \subset R^m$ be a bounded domain. Let $A \subset \Gamma \subset \Omega$ such that Γ is relatively closed in Ω, $H_{m-1}(\bar{A}) = 0$, and $\Gamma\backslash A$ is a C^1-smooth $m-1$ dimensional manifold. Suppose $\Omega\backslash\Gamma = \cup_{j=1}^{\infty}\Omega_j$, the union of at most countably many domains Ω_j. For each j, we have $\partial\Omega_j \subset \partial\Omega\cup\Gamma$. We can view $\Omega\backslash\Gamma$ as domains Ω_j obtained by cutting apart along Γ and $\Gamma\backslash A$ as the union of two copies of $\Gamma\backslash A$. Let ν_j denote the outward unit normal to $\partial\Omega_j$. Then ν_j exists for any point $p \in \partial\Omega_j \cap (\Gamma\backslash A)$. At p, there is another l (l may equal j) such that $\nu_l = -\nu_j$. Let $\vec{F} \in C^1(\Omega\backslash\Gamma)$ for simplicity and $H \in C^0(\Omega\backslash\Gamma) \cap L^1_{loc}(\Omega)$. Suppose $u \in C^1(\Omega\backslash\Gamma)$ has no singular points in $\Omega\backslash\Gamma$. Let $N_j(u)$ denote the restriction of $N(u)$ on Ω_j.

Theorem 6.3. *Suppose we have the situation described above. Furthermore, suppose* $N_j(u) \in C^0(\Omega_j \cup (\partial\Omega_j \cap (\Gamma\backslash A))) \cap C^1(\Omega_j)$ *satisfies* $divN_j(u) = H$ *in* Ω_j

for any j. *Then u is a weak solution to (1.6) in Ω if and only if for each $p \in \Gamma\backslash A$, there exist j, l as described above, such that at p, there holds*

$$(N_j(u) - N_l(u)) \cdot \nu_j = (N_j(u) - N_l(u)) \cdot \nu_l = 0.$$

We should remind the reader that for $u \in W^{1,1}(\Omega)$ and $H \in L^{\infty}(\Omega)$, u is a weak solution to (1.6) if and only if u is a minimizer for (1.7) in view of Theorem 3.3.

Proof. Let $U \subset\subset \Omega$ have compact closure in Ω, and suppose that the boundary ∂U is C^1-smooth. For $\varphi \in C_0^{\infty}(\Omega)$ with support contained in U, we compute

$$\begin{aligned} (6.14)\quad \int_U N(u) \cdot \nabla\varphi + H\varphi &= \int_{U\backslash\Gamma} N(u) \cdot \nabla\varphi + H\varphi \\ &= \int_{\partial(U\backslash\Gamma)} \varphi N(u) \cdot \nu \\ &= \sum_{(j,l)} \int_{\partial\Omega_j \cap (\Gamma\backslash A) \cap U} \varphi(N_j(u) \cdot \nu_j + N_l(u) \cdot \nu_l) \\ &= \sum_{(j,l)} \int_{\partial\Omega_j \cap (\Gamma\backslash A) \cap U} \varphi(N_j(u) - N_l(u)) \cdot \nu_j. \end{aligned}$$

For the last equality we have used $\nu_l = -\nu_j$. Now observe that u is a weak solution in Ω if and only if the first term of (6.14) vanishes for any $\varphi \in C_0^{\infty}(\Omega)$ and associated U. On the other hand, this is equivalent to concluding that $(N_j(u) - N_l(u)) \cdot \nu_j = 0$ on $\Gamma\backslash A$ by (6.14).

Q.E.D.

We remark that it is possible that $u \in C^1\backslash C^2$ while $N(u) \in C^1$ in the nonsingular domain. For instance, let $u = xy + g(y)$ with $g \in C^1\backslash C^2$. Take $\vec{F} = -\vec{X}^*$. We can then compute $N(u) = (0, \pm 1)$ in the nonsingular domain defined by $2x + g'(y) \neq 0$.

We will also make a remark on deducing the second equality in (6.14). First note that at points of A with $H_{m-1}(\bar{A}) = 0$, ν may not exist. How do we deal with this? For any $\varepsilon > 0$, we can find a finite open cover $\cup_{j=1}^k D_j \supset \bar{A}$ such that $\sum_{j=1}^k H_{m-1}(\partial D_j) < \varepsilon$. By the divergence theorem we have

$$\int_{(U\backslash\Gamma)\backslash\cup_{j=1}^k D_j} N(u) \cdot \nabla\varphi + H\varphi = \int_{\partial[(U\backslash\Gamma)\backslash\cup_{j=1}^k D_j]} \varphi N(u) \cdot \nu.$$

Passing to the limit as $\varepsilon \to 0$ and observing that the integrands are bounded (since $|N(u)| = 1$), we obtain

$$\int_{U\backslash\Gamma} N(u) \cdot \nabla\varphi + H\varphi = \int_{\partial(U\backslash\Gamma)} \varphi N(u) \cdot \nu.$$

The idea of the above argument was used in [4]. We have displayed this idea in the proof of Lemma 6.1. We also used a similar argument in the proof of Theorem 5.2 in [3]. We remark that Pauls had a similar result (for $m = 2$, $\vec{F} = -\vec{X}^*$, and $H = 0$) as Theorem C in [17]. Ritoré and Rosales also obtained a similar result for C^2-smooth minimizers (for $m = 2$, $\vec{F} = -\vec{X}^*$, and $H = constant$) as Theorem 4.15 in [18].

7. EXAMPLES

We shall give examples of Lipschitz (continuous) minimizers in dimension 2.

Definition 7.1. A p-area minimizer or a p-minimizer in short is a minimizer for (1.2) with $H = 0$.

Throughout this section, we will always work on the situation that $m = 2$, $\vec{F} = -\vec{X}^*$, and $H = 0$. Recall that the integral curves of $N^{\perp}(u)$ are straight lines (see Section 4 in [3]), called the characteristic lines, segments, or rays. We call the angle between Γ (oriented) and a characteristic ray (with direction $N^{\perp}(u)$) in B^+ (B^-, respectively) touching a point $p \in \Gamma$ the incident (reflected, respectively) angle at p. Therefore geometrically (6.11) is equivalent to saying that at $p \in \Gamma \cap B$, either $N^+(u) = N^-(u)$ (see Figure 1(b)) or $N^+(u) \neq N^-(u)$ which implies

$$\text{The incident angle} = \text{The reflected angle.} \tag{7.1}$$

(see Figure 1(a)). Suppose $u \in C^2$ at a point $p \in \Gamma \cap B$ and Γ is a singular curve. Recall that if the characteristic line segments Γ_+ and Γ_- in B^+ and B^- respectively meet at p, then $\Gamma_+ \cup \{p\} \cup \Gamma_-$ must form a straight line segment according to (the proof of) Proposition 3.5 in [3]. Therefore by (7.1) (note that $N^+(u) = -N^-(u)$ at p in this situation), we can conclude that

$$\Gamma_+ \textit{ and } \Gamma_- \textit{ are perpendicular to } \Gamma \textit{ at } p \textit{ if } u \in C^2 \textit{ at } p. \tag{7.2}$$

The constraint (7.2) gives a necessary and sufficient condition for a C^2-smooth solution of (1.1) with $H = 0$ to be a p-minimizer. We can have a function $u \in C^2(\Omega)$ which satisfies the p-minimal surface equation $divN(u) = 0$ in $\Omega \backslash S(u)$, but is not a weak solution or a p-minimizer.

Example 7.1. Consider $\vec{F} = (-y, x)$ in the following $N(u)$'s.

(a) By taking $a = \cos\vartheta$, $b = \sin\vartheta$, and $g(-bx + ay) = (\cot\vartheta)\ (-bx + ay)^2$ in (1.2) of [3] for $0 < \vartheta < \frac{\pi}{2}$, we obtain $u(x, y) = -xy + y^2 \cot\vartheta$. This is a C^2 smooth solution to $divN(u) = 0$ in $R^2 \backslash S(u)$ for $\vec{F} = (-y, x)$ by a direct computation. We can easily determine the singular set $S(u) \equiv \{u_x - y = 0,\ u_y + x = 0\} = \{y = 0\}$. On the other hand, $N^{\perp}(u) = (\cos\vartheta, \sin\vartheta)$ which is not perpendicular to the x-axis $\{y = 0\}$ (see Figure 2(a)). So in view of (7.2), this u is not a p-minimizer on any bounded domain Ω containing part of the x-axis.

(b) Let $u(x, y) = -xy + y^2 \cot\vartheta$ for $y > 0$; $= -xy + y^2 \cot\eta$ for $y < 0$; $= 0$ for $y = 0$ where $0 < \vartheta, \eta < 2\pi$, $\vartheta \neq \pi$, $\eta \neq \pi$. We compute

$$\begin{aligned} N^{\perp}(u) &= \left(\frac{\cos\vartheta}{\sin\vartheta}|\sin\vartheta|, |\sin\vartheta|\right) \text{ for } y > 0; \\ N^{\perp}(u) &= \left(-\frac{\cos\eta}{\sin\eta}|\sin\eta|, -|\sin\eta|\right) \text{ for } y < 0. \end{aligned} \tag{7.3}$$

Observe that (7.1) (or (6.11)) holds if and only if $\vartheta + \eta = 2\pi$ (see Figure 2(b)) by (7.3). Therefore we conclude that u is a ($C^{1,1}$-smooth) p-minimizer on any bounded domain in R^2 if and only if $\vartheta + \eta = 2\pi$ in view of (7.1).

Example 7.2. Let $u(x, y) = xy$ for $y > 0$, and $u = 0$ for $y \leq 0$. Consider the case of $\vec{F} = (-y, x)$. Compute



(a) (b)

FIGURE 2.

$$\begin{aligned} N^{\perp}(u) &= (1,0) \text{ for } x>0, y>0;\ N^{\perp}(u)=(-1,0) \text{ for } x<0, y>0. \\ N^{\perp}(u) &= \frac{(x,y)}{\sqrt{x^2+y^2}} \text{ for } y<0 \end{aligned}$$

(see Figure 3). Observe that the positive y-axis $\{x = 0,\ y > 0\}$ is a singular curve where (7.2) holds true. Also on the x-axis $\{y = 0\}$ except the origin, $N^{\perp}(u)$ is continuous and hence (6.11) holds true (note that the x-axis is not a singular curve, but is a curve where u is not C^1 smooth). Applying Theorem 6.3 with $\Gamma = \{x = 0,\ y > 0\} \cup \{y = 0\}$, we conclude that u is a (Lipschitz) p-minimizer on any bounded domain $\Omega \subset R^2$.



FIGURE 3.

We remark that it is not possible to construct a Lipschitz p-minimizer having a loop consisting of characteristic lines (see Figure 4 for an example). Indeed, by contradiction, suppose that the loop consists of three characteristic lines γ_1, γ_2, and γ_3 as indicated in Figure 4. Let Δ denote the region surrounded by γ_1, γ_2, and γ_3. We integrate the contact form $\Theta \equiv du + xdy - ydx$ over the loop as follows:

$$\begin{aligned} 0 &= \int_{\gamma_1 \cup \gamma_2 \cup \gamma_3} \Theta \ (\gamma_1, \gamma_2, \text{ and } \gamma_3 \text{ being Legendrian}) \\ &= \int_{\Delta} d\Theta \ \ (\text{Stokes' Theorem}) \\ &= 2\int_{\Delta} dx \wedge dy = 2 \ \text{Area}(\Delta) \neq 0. \end{aligned}$$

This contradiction confirms our claim.



Figure 4.

Example 7.3. There can be two distinct C^2-smooth p-minimal graphs (i.e., satisfying (1.1) on nonsingular domain) having the same boundary value and the same p-area, but both of them are not p-minimizers. Consider $u = x^2 + xy$, $v = xy + 1 - y^2$ (first given in [16]). We can easily verify that u and v satisfy (1.1) with $H = 0$ on their respective nonsingular domains and have the same value on the unit circle in the xy - plane. But they do not satisfy (7.2). So by Proposition 6.2 or Theorem 6.3, neither of them can be a p-minimizer. Compute the p-area (see (1.2)) of u and v over the unit disc Δ as follows:

$$\begin{aligned} \mathcal{X}(u) &= \int_{\Delta} \sqrt{8}|x| dxdy = \frac{8\sqrt{2}}{3}, \\ \mathcal{X}(v) &= \int_{\Delta} 2|x-y| dxdy = \frac{8\sqrt{2}}{3}. \end{aligned}$$

So they have the same p-area. By the uniqueness of p-minimizers (see Theorem B), we also conclude that neither u nor v can be the p-minimizer.

We are going to describe what the (unique) p-minimizer looks like on Δ with the boundary value (or curve) $\rho(\theta) \equiv \cos^2\theta + \cos\theta \sin\theta$ (θ is the standard angle parameter for $\partial\Delta$). Let (α, β, γ) be a point of a line segment $\tilde{L} \subset \bar{\Delta} \times R$ meeting the boundary curve with the projection $L \subset \bar{\Delta}$ passing through the origin. Suppose θ' is the angle between the positive x-axis and part of L, lying in the upper half plane. Then we have

$$\begin{aligned} \alpha &= t\cos\theta', \ \beta = t\sin\theta' \\ \gamma &= \cos^2\theta' + \cos\theta' \sin\theta' \end{aligned} \tag{7.4}$$

for $-1 \leq t \leq 1$ (note that $\rho(\pi+\theta) = \rho(\theta)$). Suppose the contact plane passing through $(\alpha,\ \beta,\ \gamma)$ intersects the boundary curve ρ at $(\cos\theta,\ \sin\theta,\ \rho(\theta))$. Then we have the following relation:

$$\rho(\theta) - \rho(\theta') + t\sin(\theta - \theta') = 0 \tag{7.5}$$

by observing that $z - \gamma + x(y - \beta) - y(x - \alpha) = 0$ is the equation for such a contact plane in R^3 with coordinates (x, y, z). By elementary trigonometry for the above specific ρ, we can reduce (7.5) to

$$\frac{\sqrt{2}}{2}[\sin(2\theta + \frac{\pi}{4}) - \sin(2\theta' + \frac{\pi}{4})] + t\sin(\theta - \theta') = 0. \tag{7.6}$$

The idea is to choose θ' such that $\sin(2\theta' + \frac{\pi}{4}) = 0$. Then we solve (7.6) for θ (perhaps we have multiple solutions). Keeping $\tilde{L}$ or L associated to θ' as the singular set in mind, we connect $(\alpha,\ \beta,\ \gamma) \in \tilde{L}$ to a point of the boundary curve, associated to θ, by a line segment. Since these line segments are Legendrian, their union forms a Legendrian ruled surface, hence a p-minimal surface ([3]). Moreover, if two characteristic lines (i.e., above Legendrian lines projected to the xy-plane) meet at a point of $\tilde{L}$, condition (7.1) holds. So in this way we can construct the p-minimizer by Proposition 6.2 or Theorem 6.3. We give more details below.

First solving $\sin(2\theta' + \frac{\pi}{4}) = 0$ gives $\theta' = (n - \frac{1}{4})\frac{\pi}{2}$ where n is an integer. There are two such θ''s modulo an integral multiple of π, namely $\theta' = \frac{3}{8}\pi$ and $\theta' = \frac{7}{8}\pi$. We take $\theta' = \frac{3}{8}\pi$ (it turns out that $\theta' = \frac{7}{8}\pi$ won't give rise to a p-minimal graph in the following argument). So (7.6) is reduced to

$$\frac{\sqrt{2}}{2}\sin 2(\theta - \frac{3}{8}\pi) = t\sin(\theta - \frac{3}{8}\pi). \tag{7.7}$$

(note that for $\theta' = \frac{7}{8}\pi$ we have $-t$ instead of t in (7.7)). By the double angle formula, we deduce from (7.7) that

$$(a)\ \cos(\theta - \frac{3}{8}\pi) = \frac{t}{\sqrt{2}}\ ;(b)\ \sin(\theta - \frac{3}{8}\pi) = 0. \tag{7.8}$$

The solutions to (b) of (7.8) are $\frac{3}{8}\pi\ + n\pi$ for any integer n, which we ignore. We have two solutions θ_1, θ_2 (modulo an integral multiple of 2π) to (a) of (7.8) for a given t with the relation

$$\theta_1 - \frac{3}{8}\pi = \frac{3}{8}\pi - \theta_2. \tag{7.9}$$

When t runs from -1 to 1, θ_1 runs from $\frac{9}{8}\pi$ to $\frac{5}{8}\pi$ clockwise while θ_2 runs from $-\frac{3}{8}\pi$ to $\frac{1}{8}\pi$ counterclockwise (See Figure 5 below).

Denote the line segments between $(\alpha,\ \beta) \in L$ $(\theta' = \frac{3}{8}\pi)$ and the boundary point $(\cos\theta_j,\ \sin\theta_j)$, $j = 1,\ 2$, by Γ_t^1, Γ_t^2, respectively. Γ_t^1 and Γ_t^2 are the xy-plane projections of two Legendrian lines $\tilde{\Gamma}_t^1$, $\tilde{\Gamma}_t^2$ connecting $(\alpha,\ \beta,\ \gamma) \in \tilde{L}$ to $(\cos\theta_j,\ \sin\theta_j,\ \rho(\theta_j))$, $j = 1,\ 2$, respectively. We will define a graph $\check{u}$ over $\bar{\Delta}$, whose restriction to the region $\Omega \equiv L \cup (\cup_{j=1,2;-1\leq t\leq 1}\Gamma_t^j)$ is $\tilde{L} \cup (\cup_{j=1,2;-1\leq t\leq 1}\tilde{\Gamma}_t^j)$. We can parametrize $\tilde{\Gamma}_t^2$, say, in the following form (see (4.9) in[3]):



Figure 5.

$$\begin{aligned} x &= s(\sin\eta(t)) + \alpha(t) \\ y &= -s(\cos\eta(t)) + \beta(t) \\ z &= s[\beta(t)\sin\eta(t) + \alpha(t)\cos\eta(t)] + \gamma(t). \end{aligned}$$

Here $\eta(t) = \frac{\pi}{2} + \theta_2(t) - \delta(t)$ in which $\cos(\theta_2(t) - \frac{3}{8}\pi) = \frac{t}{\sqrt{2}}$ (see (a) of (7.8)) and $\tan\delta(t) = t\sqrt{1-t^2/2}/(1-t^2/\sqrt{2})$ by elementary plane geometry (we leave the details to the reader). On the other hand, (7.1) holds along L due to (7.9). Therefore $\breve{u} \in C^{1,1}$ is a weak solution to (1.1) with $H = 0$ over the region Ω. The remaining domain $\bar{\Delta}\backslash\Omega$ consists of four small fan-shaped regions (see Figure 5). For each of such regions, we can connect two points on the boundary curve, indicated by θ' and θ which are related by (7.6) with $t = 1$. Thus we obtain a family of Legendrian line segments whose lengths are getting smaller when both θ' and θ tend to some critical value (e.g., for the fan-shaped region between $\frac{1}{8}\pi$ and $\frac{3}{8}\pi$, the critical value is $\frac{1}{4}\pi$ by solving $\frac{d\theta}{d\theta'} = 0$). These Legendrian line segments form a portion of the graph $\breve{u}$ over $\bar{\Delta}\backslash\Omega$. So $\breve{u}$ is a C^2-smooth p-minimal graph over $\bar{\Delta}\backslash\Omega$. Altogether $\breve{u} \in C^{1,1}(\bar{\Delta})$ is the (unique) p-minimizer by Theorem 3.3.

8. Appendix: uniqueness of solutions to (4.1)

The existence of solutions to (4.1) is asserted in Theorem 4.5. In this section we are going to prove the uniqueness. In fact we can obtain more general results. First we define

$$N_\varepsilon(u) \equiv \frac{\nabla u + \vec{F}}{\sqrt{\varepsilon^2 + |\nabla u + \vec{F}|^2}}. \tag{8.1}$$

Let $\vec{\alpha} \equiv (\varepsilon, \nabla u + \vec{F}) \in R \times R^m = R^{m+1}$. Denote $|\vec{\alpha}|$ as α. Similarly let $\vec{\beta} \equiv (\varepsilon, \nabla v + \vec{F})$ and $\beta \equiv |\vec{\beta}|$.

Lemma 8.1. *Let* u, $v \in W^1(\Omega)$ *where* $\Omega \subset R^m$ $(m \geq 1)$ *is an arbitrary domain. Then*

$$(N_\varepsilon(u) - N_\varepsilon(v)) \cdot (\nabla u - \nabla v) \geq \frac{\alpha+\beta}{2} \mid N_\varepsilon(u) - N_\varepsilon(v) \mid^2. \tag{8.2}$$

Moreover, the equality holds for $\varepsilon = 0$. *When* $\varepsilon > 0$, $(N_\varepsilon(u) - N_\varepsilon(v)) \cdot (\nabla u - \nabla v) = 0$ *if and only if* $\nabla u = \nabla v$.

Proof. We compute

$$\begin{aligned} & (N_\varepsilon(u) - N_\varepsilon(v)) \cdot (\nabla u - \nabla v) \\ = \;& (\frac{\nabla u + \vec{F}}{\alpha} - \frac{\nabla v + \vec{F}}{\beta}) \cdot \{(\nabla u + \vec{F}) - (\nabla v + \vec{F})\} \\ = \;& \{\frac{(\varepsilon, \nabla u + \vec{F})}{\alpha} - \frac{(\varepsilon, \nabla v + \vec{F})}{\beta}\} \cdot \{(\varepsilon, \nabla u + \vec{F}) - (\varepsilon, \nabla v + \vec{F})\} \\ = \;& \{\frac{\vec{\alpha}}{\alpha} - \frac{\vec{\beta}}{\beta}\} \cdot \{\vec{\alpha} - \vec{\beta}\} = (\alpha+\beta)(1-\cos\theta) \end{aligned} \tag{8.3}$$

where $\vec{\alpha} \cdot \vec{\beta} = \alpha\beta\cos\theta$. On the other hand, we can estimate

$$\begin{aligned} & \mid N_\varepsilon(u) - N_\varepsilon(v) \mid^2 \\ \leq \;& \mid \frac{\vec{\alpha}}{\alpha} - \frac{\vec{\beta}}{\beta} \mid^2 = 2(1-\cos\theta). \end{aligned} \tag{8.4}$$

Now (8.2) follows from (8.3) and (8.4). Observing that the equality in (8.4) holds for $\varepsilon = 0$, we obtain the equality in (8.2) for $\varepsilon = 0$. Suppose $(N_\varepsilon(u) - N_\varepsilon(v)) \cdot (\nabla u - \nabla v) = 0$. By (8.2) we have $N_\varepsilon(u) = N_\varepsilon(v)$. Taking the modulus of this equality gives $|\nabla u + \vec{F}| = |\nabla v + \vec{F}|$ if $\varepsilon > 0$. It follows that $\nabla u = \nabla v$.

Q.E.D.

We remark that the equality in (8.2) for $\varepsilon = 0$ has been obtained as Lemma 5.1′ in [3]. Recall $Q_\varepsilon u \equiv div N_\varepsilon(u)$ (see (4.1), (8.1)). Note that for $\vec{F} = 0$, $\varepsilon = 1$, $Q_\varepsilon u$ is the Riemannian mean curvature of the graph defined by u. In this case, the above inequality has been obtained in [14], [10], and [5] independently.

Definition 8.1. Let $\Omega \subset R^m$ be a bounded domain and $\varepsilon > 0$. Suppose u, $v \in W^1(\Omega)$ and $\vec{F}$ is measurable. We say $Q_\varepsilon u - Q_\varepsilon v \geq 0$ (≤ 0, respectively) weakly if for any $\varphi \in C_0^1(\Omega)$, $\varphi \geq 0$, there holds

$$\int_\Omega (N_\varepsilon(u) - N_\varepsilon(v)) \cdot \nabla\varphi \leq 0 \text{ (} \geq 0, \text{ respectively).} \tag{8.5}$$

Note that $N_\varepsilon(u)$ and $N_\varepsilon(v)$ are integrable since they are bounded by 1. We have the following comparison principle for Q_ε.

Theorem 8.2. *Let* $\Omega \subset R^m$ *be a bounded domain and* $\varepsilon > 0$. *Suppose* u, $v \in C^1(\Omega) \cap C^0(\bar{\Omega})$ *satisfy* $Q_\varepsilon u - Q_\varepsilon v \geq 0$ (≤ 0, *respectively*) *weakly and* $u - v \leq 0$ (≥ 0, *respectively*) *on* $\partial\Omega$. *Then* $u - v \leq 0$ (≥ 0, *respectively*) *in* Ω.

Proof. Given $a > 0$, we choose a function $f_a \in C^1(R)$ with the property that $f_a \equiv 0$ in $(-\infty, a]$, $f_a > 0$, and $f_a' > 0$ in (a, ∞). Observe that $f_a(u-v) \in C_0^1(\Omega)$ (i.e., $f_a(u-v) \in C^1$ and has compact support in Ω) by the assumption $u - v \leq 0$ on $\partial\Omega$. It follows from (8.5) that

$$\begin{aligned} 0 &\geq \int_\Omega (N_\varepsilon(u) - N_\varepsilon(v)) \cdot \nabla(f_a(u-v)) \\ &= \int_{\{u-v>a\}} (N_\varepsilon(u) - N_\varepsilon(v)) \cdot f_a'(u-v)(\nabla u - \nabla v) \\ &\geq 0 \ \text{ (by (8.2))}. \end{aligned} \tag{8.6}$$

Therefore we have $(\nabla u - \nabla v) \cdot (N_\varepsilon(u) - N_\varepsilon(v)) = 0$ in $\{u - v > a\}$ since $f_a'(u-v) > 0$ and $(N_\varepsilon(u) - N_\varepsilon(v)) \cdot (\nabla u - \nabla v) \geq 0$ in (8.6). It follows that $\nabla u = \nabla v$ in $\{u - v > a\}$ by Lemma 8.1. Thus we obtain $u - v \equiv a$ in $\{u - v > a\}$, a contradiction. So $\{u - v > a\}$ is empty. Since $a > 0$ is arbitrary, we conclude that $\{u - v > 0\}$ is empty. So $u - v \leq 0$ in Ω.

Q.E.D.

We remark that basically the above result can be deduced from Theorem 10.7 in [9].

Corollary 8.3. *Let $\Omega \subset R^m$ be a bounded domain. Let $\varepsilon > 0$. Suppose $u, v \in C^2(\Omega) \cap C^0(\bar{\Omega})$ and $\vec{F} \in C^1(\Omega)$ satisfy $Q_\varepsilon u = Q_\varepsilon v$ in Ω and $u = v$ on $\partial\Omega$. Then $u \equiv v$ in Ω.*

Theorem 8.4. *Let $\Omega \subset R^m$ be a bounded domain and $\varepsilon > 0$. Suppose $u, v \in W^{1,1}(\Omega)$ satisfy $Q_\varepsilon u - Q_\varepsilon v \geq 0$ (≤ 0, respectively) weakly and $(u - v)^+$ ($(u - v)^-$, respectively) $\in W_0^{1,1}(\Omega)$. Then $u - v \leq 0$ (≥ 0, respectively) in Ω.*

Proof. First we observe that (8.5) still holds for $\varphi \in W_0^{1,1}(\Omega)$, $\varphi \geq 0$. It follows that

$$\begin{aligned} 0 &\geq \int_\Omega (N_\varepsilon(u) - N_\varepsilon(v)) \cdot \nabla(u - v)^+ \\ &= \int_{\{u-v>0\}} (N_\varepsilon(u) - N_\varepsilon(v)) \cdot \nabla(u - v) \\ &\geq 0 \end{aligned} \tag{8.7}$$

by (8.2). Therefore if $u - v > 0$, $\nabla(u - v)^+ = \nabla(u - v) = 0$ by Lemma 7.6 in [9], (8.7), and Lemma 8.1. Also if $u - v \leq 0$, $\nabla(u - v)^+ = 0$ by Lemma 7.6 in [9]. Altogether we have shown that $\nabla(u - v)^+ = 0$ in Ω. Now applying the Sobolev inequality to $(u - v)^+ \in W_0^{1,1}(\Omega)$, we obtain $(u - v)^+ = 0$ in Ω. That is, $u - v \leq 0$ in Ω.

Q.E.D.

We remark that the proof of Theorem 8.4 is based on the idea of the proof of Theorem 8.1 in [9].

REFERENCES

[1] Balogh, Z. M., Size of characteristic sets and functions with prescribed gradient, J. reine angew. Math., 564 (2003) 63-83.

[2] Cheng, J.-H. and Hwang, J.-F., Properly embedded and immersed minimal surfaces in the Heisenberg group, Bull. Aus. Math. Soc., 70 (2004) 507-520.

[3] Cheng, J.-H., Hwang, J.-F., Malchiodi, A., and Yang, P., Minimal surfaces in pseudohermitian geometry, Annali della Scuola Normale Superiore di Pisa, Classe di Scienze (5), 4 (2005) 129-177.

[4] Concus, P. and Finn, R., On capillary free surfaces in the absence of gravity, Acta Math., 132 (1974) 177-198.

[5] Collin, P. and Krust, R., Le Problème de Dirichlet pour l'équation des surfaces minimales sur des domaines non bornès, Bull. Soc. Math. France, 119 (1991) 443-462.

[6] Franchi, B., Serapioni, R., and Serra Cassano, F., Rectifiability and perimeter in the Heisenberg group, Math. Ann. 321 (2001) 479-531.

[7] Garofalo, N. and Nhieu, D.-M., Isoperimetric and Sobolev inequalities for Carnot-Caratheodory spaces and the existence of minimal surfaces, Comm. Pure Appl. Math., 49 (1996) 1081-1144.

[8] Garofalo, N. and Pauls, S. D., The Bernstein problem in the Heisenberg group, preprint, 2005

[9] Gilbarg, D. and Trudinger, N. S., Elliptic partial differential equations of second order, 2nd ed., G.M.W. 224, Springer-Verlag, 1983.

[10] Hwang, J.-F., Comparison principles and Liouville theorems for prescribed mean curvature equation in unbounded domains, Ann. Scuola Norm. Sup. Pisa, 15 (1988) 341-355.

[11] Juutinen, P. and Lindqvist, P., A theorem of Radó's type for the solutions of a quasi-linear equation, Math. Res. Lett., 11 (2004) 31-34.

[12] Juutinen, P., P-harmonic approximation of functions of least gradient, to appear in Indiana Univ. Math. J..

[13] Lee, J. M., The Fefferman metric and pseudohermitian invariants, Trans. Amer. Math. Soc., 296 (1986) 411-429.

[14] Miklyukov, V. M., On a new approach to Bernstein's theorem and related questions for equations of minimal surface type, Mat. Sb., 108(150) (1979) 268-289; English transl. in Math. USSR Sb., 36 (1980) 251-271.

[15] Morrey, C. B., Multiple integrals in the calculus of variations, GMW 130, Springer-Verlag New York Inc. 1966.

[16] Pauls, S. D., Minimal surfaces in the Heisenberg group, Geometric Dedicata, 104 (2004) 201-231.

[17] Pauls, S. D., H-minimal graphs of low regularity in H^1, Comment. Math. Helv., 81 (2006) 337-384.

[18] Ritoré, M. and Rosales, C., Area-stationary surfaces in the Heisenberg group H^1, arXiv: math.DG/0512547 v1.

[19] Sternberg, P., Williams, G., and Ziemer, W. P., Existence, uniqueness, and regularity for functions of least gradient, J. reine angew. Math., 430 (1992) 35-60.

(Jih-Hsin Cheng and Jenn-Fang Hwang) INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, TAIPEI, TAIWAN, R.O.C.
E-mail address, Jih-Hsin Cheng: `cheng@math.sinica.edu.tw`
URL: `http://www.math.sinica.edu.tw`

E-mail address, Jenn-Fang Hwang: `majfh@math.sinica.edu.tw`
URL: `http://www.math.sinica.edu.tw`

(Paul Yang) DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544, U.S.A.
E-mail address: `yang@Math.Princeton.EDU`


y
x
(0,1)
$(0,\frac{1}{2})$
(0,0)